

B9

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 199 301 B2

(12)

NEW EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the opposition decision:
09.07.1997 Bulletin 1997/28

(51) Int. Cl.⁶: C07K 14/16, C12N 15/00,
C12P 21/02, C12P 21/00,
G01N 33/569, A61K 39/21

(45) Mention of the grant of the patent:
29.12.1993 Bulletin 1993/52

(21) Application number: 86105371.8

(22) Date of filing: 18.04.1986

(54) **Recombinant acquired immune deficiency syndrome (AIDS) viral envelope protein fragments and method of testing for AIDS**

Rekombinantes Viren-Überzugsprotein assoziiert mit "Acquired Immune Deficiency Syndrome" (AIDS) und Verfahren zur Testung von AIDS

Protéine recombinante d'enveloppe du virus du syndrome d'immunodéficience acquise (SIDA) et procédé pour l'analyse du SIDA

(84) Designated Contracting States:
AT BE CH DE FR GB IT LI NL SE

(30) Priority: 19.04.1985 US 725021

(43) Date of publication of application:
29.10.1986 Bulletin 1986/44

(73) Proprietors:
• F. HOFFMANN-LA ROCHE AG
4002 Basel (CH)
• THE GOVERNMENT OF THE UNITED STATES OF
AMERICA as represented by THE SECRETARY
of the DEPARTMENT OF HEALTH AND HUMAN
SERVICES
Washington, DC 20201 (US)

(72) Inventors:
• Crowl, Robert Mitchell
Cedar Grove, N.J. 07009 (US)
• Gallo, Robert Charles
Bethesda Maryland 20817 (US)
• Reddy, Eragam Premkumar
Montclair, N.J. 07042 (US)
• Shaw, George Mead
Birmingham Alabama 35243 (US)
• Wong-Staal, Flossie Yeeching
Germantown Maryland 20874 (US)

(74) Representative: Lederer, Franz, Dr. et al
Lederer, Keller & Riederer
Patentanwälte
Prinzregentenstrasse 16
80538 München (DE)

(56) References cited:
EP-A- 0 152 030 EP-A- 0 173 529
EP-A- 0 181 150 WO-A-84/04327
US-A- 4 520 113

- SCIENCE, vol. 228, no. 4695, 5th April 1985, pages 93-96; N. CHANG et al.: "Expression in Escherichia coli of open reading frame gene segments of HTLV-III"
- NATURE, vol. 313, no. 6002, 7th February 1985, pages 450-458; M.A. MUESING et al.: "Nucleic acid structure and expression of the human AIDS/lymphadenopathy retrovirus"
- NATURE, vol. 313, no. 6000, 24th January 1985, pages 277-284, London, GB; L. RATNER et al.: "Complete nucleotide sequence of the AIDS virus, HTLV-III"
- SCIENCE, vol. 226, no. 4679, 7th December 1984, pages 1165-1171; G.M. SHAW et al.: "Molecular characterization of human T-cell leukemia (lymphotropic) virus type III in the acquired immune deficiency syndrome"

EP 0 199 301 B2

Description

The present invention relates to an envelope protein fragments of an acquired immune deficiency syndrome (AIDS) virus, essentially free of other proteins, with the amino acid sequence:

5 ValTrpLysGluAla
 ThrThrThrLeuPheCysAlaSerAspAlaLysAlaTyrAspThrGluValHisAsnValTrpAlaThr
 HisAlaCysValProThrAspProAsnProGlnGluValValLeuValAsnValThrGluAsnPheAsn
 10 METrpLysAsnAspMETValGluGlnMETHisGluAspIleIleSerLeuTrpAspGlnSerLeuLys
 ProCysValLysLeuThrProLeuCysValSerLeuLysCysThrAspLeuLysAsnAspThrAsnThr
 AsnSerSerSerGlyArgMETIleMETGluLysGlyGluIleLysAsnCysSerPheAsnIleSerThr
 SerIleArgGlyLysValGlnLysGluTyrAlaPhePheTyrLysLeuAspIleIleProIleAspAsn
 AspThrThrSerTyrThrLeuThrSerCysAsnThrSerValIleThrGlnAlaCysProLysValSer
 15 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 20 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 25 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 30 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 35 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

40 CysProLysValSer
 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 45 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 50 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 55 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

5 METArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 10 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 15 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

20
 25 METTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 30 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

35
 40 METArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 45 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer.

50 It also relates to an expression vector comprising a gene coding for an envelope protein as defined above, to trans-
 formants and methods for the production of said proteins and a method for detecting the presence of AIDS antibodies
 in human blood.

Background of the Invention

55 From 1981 to date, there have been more than eight thousand (8,000) people diagnosed as having acquired
 immune deficiency syndrome (AIDS) [N.Y. Times, A-11 January 11, 1985]. AIDS has been characterized by the onset
 of severe opportunistic infections secondary to an effect on the body's immune system [Gottlieb, M.S. et al., "Pneumo-
 cystis Carinii Pneumonia and Mucosal Candidiasis in previously healthy homosexual men: evidence of a new acquired
 cellular immunodeficiency", N. Eng. J. Med. 305, 1426-1431 (1981)]. The disease has been found in male homosexu-
 als, patients receiving blood products, intravenous drug addicts, and individuals originating from Haiti and Central Africa

[Pot, P. et al., "Acquired immunodeficiency syndrome in a heterosexual population in Zaire", *Lancet* 11, 65-69 (1984)]. The causative agent was suspected to be of viral origin as the epidemiological pattern of AIDS was consistent with a transmissible disease. At least three (3) retroviruses have been isolated from cultured T-cells of several patients with AIDS, or from white blood cells of persons at risk for the disease. A novel human retrovirus called lymphadenopathy-associated virus (LAV) was discovered and its properties were consistent with its etiological role in AIDS. That virus was isolated from a patient with lymphadenopathy and hence the name [Montagnier, L. et al., "A New Human T-lymphotropic retrovirus: characterization and possible role in lymphadenopathy and acquired immune deficiency syndromes. In Human T-Cell Leukemia/Lymphoma Virus, R.C. Gallo, M. Essex and L. Gross, eds. (Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory) pp. 363-370]. Other human retroviruses, specifically two subgroups of the human T-cell leukemia/lymphoma/lymphotropic virus, types I and III have been isolated [HTLV I: Poiesz, B.J. et al., "Detection and isolation of type C retrovirus particles from fresh and cultured lymphocytes of a patient with cutaneous T-cell lymphoma", *PNAS (USA)* 77, 7415-7419 (1980); HTLV-III: Popovic, M. et al., "Detection, isolation and continuous production of cytopathic retroviruses (HTLV-III) from patients with AIDS and pre-AIDS", *Science* 224, 497-500 (1984)]. Still another virus, the AIDS-associated retrovirus (ARV), was proposed as the causative agent [Levy, J.A. et al., "Isolation of lymphocytopathic retroviruses from San Francisco patients with AIDS", *Science* 225, 840-842 (1984)]. Both the HTLV-III and ARV retroviruses display biological and sero-epidemiological properties similar to LAV [Levy J.A. et al., supra, Popovic, M. et al., supra]. As seen from the above, at least three (3) retroviruses have been postulated as the etiologic agent of AIDS: LAV; ARV; and, HTLV subtypes I and III.

LAV, HTLV III and ARV-II genomes have been molecularly cloned [Schüpbach, J. et al., "Serological analysis of a subgroup of human T-lymphotropic retroviruses (HTLV-III) associated with AIDS", *Science* 224, 503-505 (1984); Alizon, M. et al., "Molecular Cloning of lymphadenopathy - associated virus", *Nature* 312, 757-760 (1984)]. The complete nucleotide sequence of the proviral genome of LAV, ARV and HTLV III has been determined [Ratner, L. et al., "Complete nucleotide sequence of the AIDS virus, HTLV III", *Nature* 313, 277-284 (1985); Sanchez-Pescador, R. et al., "Nucleotide sequence and expression of an AIDS-associated retrovirus (ARV-2)", *Science* 227, 484-492 (1985); Wain-Hobson, S. et al., "Nucleotide sequence of the AIDS virus, LAV", *Cell* 40, 9-17 (1985)].

Shaw et al., *Science* 226, 1165-1171 (1984), describes the molecular cloning and analysis of the full-length HTLV-III proviral genome comparing various DNA-clones.

Another analysis of the HTLV-III genome is shown by Muesing et al., *Nature* 313, 450-458 (1985).

Chang et al., *Science* 228, 93-96 (1985), describes the expression of small DNA fragments fused to DNA sequences encoding the λ CI protein and β -galactosidase resulting in unpurified env polypeptides fused to the λ CI protein at their amino termini and to β -galactosidase at their carboxyl termini.

One reason for the difficulty in determining the etiologic agent of AIDS was due to the reactivity of various retroviral antigens with serum samples from AIDS patients. For example, serum samples from AIDS patients have been shown to react with antigens of HTLV I and HTLV III [HTLV-I: Essex, M. et al., "Antibodies to Cell Membrane Antigens Associated with Human T-Cell Leukemia Virus in Patients with AIDS", *Science* 220, 859-862 (1983); HTLV-III: Sarngadharan, M.G. et al., "Antibodies Reactive With Human T-Lymphotropic Retroviruses (HTLV-III) in the Serum of Patients With AIDS", *Science* 224, 506-508 (1984)]. Envelope gene products of HTLV demonstrated antigenicities cross-reactive with antibodies in sera from adult T-cell leukemia patients [Kiyokawa, T. et al., "Envelope proteins of human T-cell leukemia virus: Expression in *Escherichia coli* and its application to studies of env gene functions", *PNAS (USA)* 81, 6202-6206 (1984)]. Adult T-cell leukemias (ATL) differ from acquired immune deficiency syndrome (AIDS) in that HTLV-I causes T-cell malignancies, that is uncontrolled growth of T-cell. In AIDS rather than cell growth there is cell death. In fact this cytopathic characteristic of HTLV III was critical to determining ultimately the specific retroviral origin of the disease. Thus the etiologic agent of AIDS was isolated by use of immortalized human neoplastic T cell lines (HT) infected with the cytopathic retrovirus characteristic of AIDS, isolated from AIDS afflicted patients. Seroepidemiological assays using this virus showed a complete correlation between AIDS and the presence of antibodies to HTLV III antigens [Sarngadharan, M.G. et al., supra; Schupbach, J. et al., supra]. In addition, nearly 85% of patients with lymphadenopathy syndrome and a significant proportion of asymptomatic homosexual men in AIDS endemic areas were also found to carry circulating antibodies to HTLV III. Taken together, all these data indicate HTLV III to be the etiologic agent for AIDS.

Until the successful culturing of AIDS virus using H-9 cell line [PCT application, publication no. WO 85/04897] the env AIDS protein of the AIDS virus had not been isolated, characterized or synthesized. This in major part is due to the fact that the virus is cytopathic and thus isolation of the virus was not possible [Popovic, M. et al., supra]. Once the human T-cell line resistant to the cytopathic effects of the virus was discovered, a molecular clone of proviral DNA could be achieved.

The need for a sensitive and rapid method for the diagnosis of AIDS in human blood and its prevention by vaccination is very great. Virtually all the assays/tests presently available are fraught with errors. In fact the Center for Disease Control (CDC) has indicated that presently available tests be used solely for screening units of blood for antibody to HTLV III. The CDC went further by stating that the presently available ELISA tests can not be used for general screening of high risk populations or as a diagnostic test for AIDS [Federal Register 50(48), 9909, March 12, 1985]. The errors have been traced to the failure to use a specific antigenic protein of the etiologic agent for AIDS. The previously used

proteins were derived from a viral lysate. Since the lysate is made from human cells infected with the virus, i.e. the cells used to grow the virus, the lysate will contain human proteins as well as viral proteins. Thus preparation of a pure antigen of viral protein is very difficult. The antigen used produced both false positive and false negative results [Budiansky, S., "AIDS Screening, False Test Results Raise Doubts", Nature 312, 583(1984)]. The errors caused by the use of such lysate proteins/peptides can be avoided by using a composition for binding AIDS antibodies which is substantially free of the non-AIDS specific proteins. Compositions that are substantially pure AIDS envelope protein can be used as antigens.

The AIDS envelope protein of the instant invention has been established to have conserved epitopes which permit its use to screen for, diagnose and/or prevent by vaccination the infection by AIDS virus. The instant invention demonstrates that the envelope protein with its conserved epitopes includes all the variants which have been claimed as the sole etiologic agent.

The envelope AIDS protein of the present invention may be produced by conventionally known methods. The processes by which the novel protein may be produced can be divided into three groups: (1) chemical synthesis; (2) preparation of a gene prepared by chemical synthesis which is inserted into a host and a protein is produced by the host; and (3) a corresponding gene obtained biotechnically is inserted into a host and a protein is produced by the host.

In one embodiment of this invention, recombinant DNA techniques are utilized by which env AIDS DNA from a natural source is introduced into a cell to produce the env AIDS protein. One method of obtaining DNA which encodes env AIDS is to read the genetic code in reverse and synthesize an oligodeoxynucleotide which should encode the env AIDS amino acid sequence. As the env protein has not been isolated or characterized this approach cannot be pursued.

Alternatively gene expression can be obtained using recombinant DNA technology if DNA isolated from natural sources is used instead of synthetic DNA.

Summary of the Invention

This invention is directed to the engineering of HTLV III env gene into suitable expression vectors; transformation of host organisms with such expression vectors; and production of envelope AIDS protein (env AIDS) by culture of such transformed cells. Another aspect of the present invention relates to the isolation and use of the resulting recombinant env AIDS protein.

Another aspect of the present invention is the identification and determination of the proviral DNA sequence. More specifically, this aspect of the invention relates to determination and comparison of the proviral nucleotide sequence of the envelope genes of the purported etiologic agent of AIDS i.e. lymphadenopathy-associated virus (LAV), AIDS-associated retrovirus (ARV) and the human T-cell leukemia/lymphoma/lymphotropic virus type III (HTLV III).

A further aspect of this invention relates to a diagnostic method for testing human blood for the presence of antibodies to the env AIDS protein. This aspect of the invention overcomes the problems of all previously used blood tests for AIDS. One of the problems is the use of compositions to bind AIDS antibody which contain proteins or peptides which were not derived solely from the AIDS etiologic agent. A composition using homogeneous envelope AIDS protein of this invention overcomes the nonspecificity of the prior tests or assays. Yet another aspect of this invention is a diagnostic method for detecting and/or determining the presence of the antigen in human blood.

Another aspect of this invention is to use the env AIDS proteins of the instant invention as antigens suitable for providing protective immunity against AIDS when incorporated into a vaccine.

Brief Description of the Drawings

Fig. 1. The nucleotide sequence of the envelope gene of the HTLV-III proviral genome (HXB-3).

Fig. 2. Comparison of the amino acid sequence of the env protein of the five purported etiologic agents of AIDS. Amino acid sequences are aligned to give maximum homology.

Fig. 3. Construction of the pEV/env44-640 expression plasmids. The upper left panel shows a simplified restriction site map of the 3.15 Kb EcoRI-XhoI segment of the HTLV-III genome which contains the env coding region (cross-hatched arrow). The right panel shows the structure and pertinent sequences of the pEV-vrf plasmids. The solid black region represents the synthetic ribosome binding site sequences upstream of the ATG initiation codon (overlined). See Example 2 for a detailed description of the env expression plasmid constructions.

Fig. 4. Western blot analysis of env coded antigens produced in E. coli. Total bacterial proteins were resolved by SDS-PAGE, electro-blotted onto a nitrocellulose filter, and env encoded proteins were detected by reacting with human sera as described in Example 5: a) negative control, cells containing pJCL-E30 (p21T) induced at 42° C for 2 hours; b) uninduced control, cells containing pEV3/env44-640 maintained at 30° C; c) pEV3/env44-640; d) pEV1/env44-640; and e) pEV3/env205-640 induced at 42° C for 2 hours.

Fig. 5. Recognition of bacterially synthesized HTLV-III env gene products by antibodies in AIDS patient sera. Bacterial lysates containing recombinant env proteins were subjected to Western blot analysis as described in Example 5.

Individual strips were then incubated with a 1000-fold dilution of individual sera followed by treatment with ^{125}I -labeled protein A. (upper part) Serum samples were from the following donors: (lane 1) normal healthy donor; (lanes 2-18) AIDS patient sera collected from the West Coast of the USA. (Lower part) Serum samples were taken from the following donors: (lane 1) donor found to be HTLV-1(+) by Elisa using disrupted virus; (lanes 4, 5, 11 and 15) healthy, normal donors; (lanes 2, 3, 6, 8, 10, 12, 13, 14, 16, 17 and 18) AIDS patient sera from the East Coast of the USA.

Fig. 6A. The amino acid sequence of the AIDS envelope protein.

Fig. 6B. The amino acid distribution of the AIDS envelope protein.

Fig. 7. Construction of the expression vector pRC23. The Shine-Dalgarno sequence (SD) is overlined and the location of the synthetic ribosome binding site sequence in the plasmid is represented by the solid black segment. The plasmid contains the entire sequence of pBR322 and thus confers resistance to both ampicillin (amp^R) and tetracycline (tet^R).

Fig. 8. Construction of the pEV-vrf vectors. The synthetic oligonucleotides for each plasmid which were placed downstream of the SD sequence in pRC23 are shown with the locations of the restriction enzyme cleavage sites. The ATG initiation codon is overlined, and the placement of the additional A-T base pairs is designated by the rectangle. The plasmids confer resistance to ampicillin only.

Detailed Description of the Invention

In the description the following terms are employed:

Nucleotide: A monomeric unit of DNA consisting of a sugar moiety (pentose), a phosphate, and either a purine or pyrimidine base (nitrogenous heterocyclic). The base is linked to the sugar moiety via the glycosidic carbon (1' carbon of the pentose). That combination of a base and a sugar is called a nucleoside. Each nucleotide is characterized by its base. The four DNA bases are adenine ("A"), guanine ("G"), cytosine ("C") and thymine ("T").

DNA Sequence: A linear array of nucleotides connected one to the other by phosphodiester bonds between the 3' and 5' carbons of adjacent pentoses.

Codon: A DNA sequence of three nucleotides (a triplet) which encodes through mRNA an amino acid, a translation start signal or a translation termination signal. For example, the nucleotide triplets TTA, TTG, CTT, CTC, CTA and CTG encode for the amino acid leucine ("Leu"). TAG, TAA and TGA are translation stop signals and ATG is a translation start signal.

Reading Frame: The grouping of codons during translation of mRNA into amino acid sequences. During translation the proper reading frame must be maintained. For example, the sequence GCTGGTTGTAAG may be translated in three reading frames or phases, each of which affords a different amino acid sequence:

GCT GGT TGT AAG=Ala-Gly-Cys-Lys

G CTG GTT GTA AG=Leu-Val-Val

GC TGG TTG TAA G=Trp-Leu-(STOP)

Polypeptide: A linear array of amino acids connected one to the other by peptide bonds between the α -amino and carboxy groups of adjacent amino acids.

Genome: The entire DNA of a cell or a virus. It includes *inter alia* the structural genes coding for the polypeptides of the substance, as well as operator, promoter and ribosome binding and interaction sequences, including sequences such as the Shine-Dalgarno sequences.

Structural Gene: A DNA sequence which encodes through its template or messenger RNA ("mRNA") a sequence of amino acids characteristic of a specific polypeptide.

Transcription: The process of producing mRNA from a structural gene.

Translation: The process of producing a polypeptide from mRNA.

Expression: The process undergone by a structural gene to produce a polypeptide. It is a combination of transcription and translation.

Plasmid: A circular double-stranded DNA molecule that is not a part of the main chromosome of an organism containing genes that convey resistance to specific antibiotics. When the plasmid is placed within a unicellular organism, the characteristics of that organism may be changed or transformed as a result of the DNA of the plasmid. For example, a plasmid carrying the gene for tetracycline resistance (Tet^R) transforms a cell previously sensitive to tetracycline into one which is resistant to it. A cell transformed by a plasmid is called a "transformant".

Cloning Vehicle: A plasmid, phage DNA or other DNA sequences which are able to replicate in a host cell, which are characterized by one or a small number of endonuclease recognition sites at which such DNA sequences may be cut in a determinable fashion without attendant loss of an essential biological function of the DNA, e.g., replication, production of coat proteins or loss of promoter or binding sites, and which contain a marker suitable for use in the identification of transformed cells, e.g., tetracycline resistance or ampicillin resistance. A cloning vehicle is often called a vector.

Cloning: The process of obtaining a population of organisms or DNA sequences derived from one such organism

or sequence by asexual reproduction.

Recombinant DNA Molecule or Hybrid DNA: A molecule consisting of segments of DNA from different genomes which have been joined end-to-end outside of living cells and have the capacity to infect some host cell and be maintained therein.

5

The nomenclature used to define the peptides or proteins is that used in accordance with conventional representation such that the amino group at the N-terminus appears to the left and the carboxyl group at the C-terminus to the right. By natural amino acid is meant one of the amino acids commonly occurring in natural proteins comprising Gly, Ala, Val, Leu, Ile, Ser, Thr, Lys, Arg, Asp, Asn, Glu, Gln, Cys, Met, Phe, Tyr, Pro, Trp and His. By Nle is meant norleucine, and by Nva is meant norvaline. Where L and D forms are possible, it is the L-form of the amino acid that is represented unless otherwise expressly indicated. In addition, amino acids have been designated by specific letters of the alphabet such that: A=Alanine; B = Aspartic Acid or Asparagine; C = Cysteine; D = Aspartic Acid; E = Glutamic Acid; F = Phenylalanine; G = Glycine; H = Histidine; I = Isoleucine; K = Lysine; L = Leucine; M = Methionine; N = Asparagine; P = Proline; Q = Glutamine; R = Arginine; S = Serine; T = Threonine; V = Valine; W = Tryptophan; Y = Tyrosine; Z =

15

Glutamine or Glutamic Acid.

In accordance with the present invention, the search for the envelope protein of the etiologic agent for acquired immune deficiency syndrome (AIDS) has led to the isolation and sequencing of the proviral gene of the AIDS virus. It has now been discovered, for what is believed to be the first time that the postulated etiologic agents of AIDS, lymphadenopathy-associated virus (LAV), AIDS-associated retrovirus (ARV) and human T-cell leukemia/lymphoma/lymphotropic virus (HTLV III) are in fact variants of the same virus. For purposes of this invention, in the specification and claims the virus causing AIDS will be referred to herein as AIDS virus. AIDS virus will be understood to include the variants which have been postulated as the causative agents of AIDS, namely LAV, ARV and HTLV III. The envelope protein of the AIDS virus (env AIDS) is a 97,200 dalton protein with 32 potential N-glycosylation sites. Nucleotide sequence analysis of the AIDS envelope gene of the putative etiologic agents of AIDS demonstrates that all the viruses are variants of the same virus. That means that there is approximately 1 to 20% divergence or variation from the sequence of the envelope gene of HTLV III and the sequences of the envelope genes of the other viruses LAV and ARV-2. The amino acid sequence of the env AIDS is set forth in Figure 6(a). The amino acid distribution is set forth in Figure 6(b).

20

25

The nucleotide sequence of the envelope gene is shown in Figure 1. The proviral DNA sequence, using methods known to one of ordinary skill in the art such as the chemical degradation method of Maxam and Gilbert of the M13 sequencing system of Messing which is a modification of the dideoxy nucleotide chain termination method of Sanger, was analyzed to determine the location of the region coding for the envelope protein. The location of an open reading frame, i.e. a long stretch of triplet codons not interrupted by a translational stop codon, for the envelope gene was determined. The open reading frame coding for the env gene is 863 amino acids and contained an ATG codon at the eighth position from the 5' end of the reading frame. The ATG codon is known to be a universal translation-initiation codon.

30

The integrated proviral genome of HTLV-III was cloned from the genomic DNA of H9 cells infected with HTLV-III [Shaw, G.M. et al., "Molecular characterization of Human T-cell leukemia (lymphotropic) virus type III in the acquired immune deficiency syndrome", Science 226, 1165-1171 (1984)]. Since the HTLV-III provirus was found to lack XbaI restriction sites, a genomic library was constructed by using XbaI digested H9/HTLV-III DNA. There are several methods available to one of ordinary skill in the art for screening the bacterial clones containing the AIDS env protein cDNA. These include, for example, RNA selection hybridization, differential hybridization with a synthetic probe or screening for clones that produce the desired protein by immunological or biological assays. From the genomic library, colonies of cells transformed with DNA that contains the HTLV III sequences were selected by hybridization screening of the library with HTLV III cDNA. The DNA insert of the hybridization-positive clone, HXB-3, was excised from the plasmid DNA and sequenced.

35

40

The predicted product of the env gene shares many features in common with the envelope gene products of other retroviruses. Thus, a hydrophobic region is seen in the middle of the protein (amino acids 519-534) which includes a processing site for the cleavage of the precursor protein into exterior and transmembrane proteins. Similarly, the amino terminal end contains a short stretch of hydrophobic amino acids (amino acids 17-37) which constitutes a potential signal sequence. The HTLV-III envelope precursor differs from the other retroviral envelope protein precursors in that it contains an additional stretch of 180 amino acids at the carboxy terminus.

45

50

Polymorphism within the Envelope Region of AIDS Virus

The recent publication of the nucleotide sequences of LAV, ARV-2 and HTLV-III [Ratner, L., et al., supra; Sanchez-Pescador, R., et al., supra; Wain-Hobson, S., et al., supra] allows a detailed comparison of these various isolates obtained from AIDS patients from different parts of the world. HTLV-III clones were isolated from AIDS patient lymphocytes obtained from the east coast of the United States, while LAV was isolated from a French man and ARV was isolated from a patient in California. A comparison of the sequence data confirms the earlier observations made using restriction enzyme site analysis which showed approximately 10% variation. The present analysis shows that the vari-

55

ous isolates show the greatest amount of conservation in the gag and pol regions while the most divergence occurs in the env region. A comparison of the five env sequences is presented in Figure 2. With respect to the envelope gene, HTLV-III and LAV are more closely related to each other than the ARV clone. Approximately 1.6% divergence was observed between the HTLV-III (HXB-3) and LAV sequence. Among the HTLV sequences, the divergence was about 1.6%. However, approximately 17% divergence was observed between HTLV-III and ARV-2 and this was more pronounced in the extracellular region of the envelope gene product (Figure 2). This high rate of divergence could be due to the geographical location from where the two isolates were derived or the time of isolation of these variants. ARV-2 was isolated from the west coast of the United States more recently. The HTLV-III isolates for which the nucleotide sequences have been determined were all obtained from the east coast of the United States a year earlier. LAV was obtained from a French patient who appears to have acquired the virus in New York about the same period. The observed differences in the sequence probably reflect divergent evolution of strains separated in time or geography or both. Within the env region, the highest level of divergence is in the extracellular portion of the protein.

Expression Vector

A wide variety of host/cloning vehicle combinations may be employed in cloning the double-stranded DNA. For example, useful cloning vehicles may consist of segments of chromosomal, nonchromosomal and synthetic DNA sequences, such as various known bacterial plasmids, e.g. plasmids from *E. coli* such as pBR322, phage DNA, and vectors derived from combinations of plasmids and phage DNAs such as plasmids which have been modified to employ phage DNA or other expression control sequences or yeast plasmids. Useful hosts may include microorganisms, mammalian cells, plant cells and the like. Among them microorganisms and mammalian cells are preferably employed. As preferable microorganisms, there may be mentioned yeast and bacteria such as *Escherichia coli*, *Bacillus subtilis*, *Bacillus stearothermophilus* and *Actinomyces*. The above-mentioned vectors and hosts may also be employed for the production of a protein from a gene obtained biologically as in the instant invention. Of course, not all host/vector combinations may be equally efficient. The particular selection of host/cloning vehicle combination may be made by those of skill in the art after due consideration of the principles set forth without departing from the scope of this invention.

Furthermore, within each specific cloning vehicle, various sites may be selected for insertion of the double-stranded DNA. These sites are usually designated by the restriction endonuclease which cuts them. For example, in pBR322 the *EcoRI* site is located just outside the gene coding for ampicillin resistance. Various sites have been employed by others in their recombinant synthetic schemes. Several sites are well recognized by those of skill in the art. It is, of course, to be understood that a cloning vehicle useful in this invention need not have a restriction endonuclease site for insertion of the chosen DNA fragment. Instead, the vehicle could be joined to the fragment by alternative means.

The vector or cloning vehicle and in particular the site chosen therein for attachment of a selected DNA fragment to form a recombinant DNA molecule is determined by a variety of factors, e.g., number of sites susceptible to a particular restriction enzyme, size of the protein to be expressed, susceptibility of the desired protein to proteolytic degradation by host cell enzymes, contamination of the protein to be expressed by host cell proteins difficult to remove during purification, expression characteristics, such as the location of start and stop codons relative to the vector sequences, and other factors recognized by those of skill in the art. The choice of a vector and an insertion site for a particular gene is determined by a balance of these factors, not all selections being equally effective for a given case.

There are several known methods of inserting DNA sequences into cloning vehicles to form recombinant DNA molecules which are equally useful in this invention. These include, for example, direct ligation, synthetic linkers, exonuclease and polymerase-linked repair reactions followed by ligation, or extension of the DNA strand with DNA polymerase and an appropriate single stranded template followed by ligation.

The cloning vehicle or vector containing the foreign gene is employed to transform a host so as to permit that host to express the protein or portion thereof for which the hybrid DNA codes. The selection of an appropriate host is also controlled by a number of factors recognized by the art. These include, for example, compatibility with the chosen vector, toxicity of proteins encoded by the hybrid plasmid, ease of recovery of the desired protein, expression characteristics, biosafety and costs. A balance of these factors must be struck with the understanding that not all hosts may be equally effective for expression of a particular recombinant DNA molecule.

A preferred embodiment of the instant invention is to express segments of the AIDS env protein in *E. coli* by inserting restriction fragments isolated from the cloned proviral genome into the versatile pEV-vrf (variable reading frame) expression plasmids (for details of construction see Example 2). These versatile pEV-vrf plasmids are derivatives of pBR322 which contain the phage lambda P_L promoter, a synthetically-derived ribosome-binding site, and convenient cloning sites (*EcoRI*, *BamHI*, *ClaI* and *HindIII*) just down-stream to the initiation codon (Figure 8). A set of three plasmids was constructed to accommodate all three translational reading frames. The P_L promoter is regulated by a temperature-sensitive *cl* repressor encoded on the compatible plasmid pRK248cits [ATCC 33766; Bernard, H.U. and Helinski, D.R., "The use of the λ phage promoter P_L to promote gene expression in hybrid plasmid cloning vehicles", *Meth. Enzymol.* 68, 482-492 (1979)]. These expression plasmids have been used to produce substantial amounts of several het-

erologous proteins in *E. coli* including v-bas p21 [Lacal, J.C. et al., "Expression of Normal and Transforming H-ras genes in *E. coli* and purification of their encoded p21 proteins", *PNAS* 81, 5305-5309 (1984)] and murine interleukin-1 [Lomedico, P.T. et al., "Cloning and Expression of Murine Interleukin-1 cDNA in *E. coli*", *Nature* 312, 458-462 (1984)].

In the present synthesis the preferred initial cloning vehicle is the bacterial plasmid pBR322 (ATCC 37017) and the preferred initial restriction endonuclease sites therein are the *EcoRI* and *HindIII* sites (Figure 3). Insertion of proviral DNA contained within the genome of H9 cells into these sites provides a large number of bacterial clones each of which contains one of the proviral DNA genes or fragments thereof present in the genome of H9 cells. Only a very few of these clones will contain the gene for env AIDS or fragments thereof.

The preferred host for initial cloning and expression of the env AIDS gene in accordance with this invention is *E. coli* MC 1061 [Casadaban, M.J. and Cohen, S.M., "Analysis of Gene Control Signals by DNA Fusion and Cloning in *E. coli*", *J. Mol. Biol.*, 138, 179-207 (1980)].

The coding sequences for amino acid residues #44 to 640 of the env protein are located downstream of the P_L promoter between the *KpnI* and *HindIII* sites on the restriction map as shown in Figure 3. Aside from the location of these convenient restriction sites, these sequences were chosen for bacterial expression experiments because they did not include the amino-terminal signal peptide as well as the hydrophobic transmembrane segment at the carboxyl end. These sequences were excluded to avoid possible toxicity problems which can occur when hydrophobic proteins are over-produced in bacterial cells. In a preferred embodiment of this invention an expression plasmid was constructed that would direct the synthesis of this segment of the env gene product (designated pEV/env 44-640), an intermediate construction was first made by inserting a 2400 bp *EcoRI*-*HindIII* fragment between the *EcoRI* and *HindIII* sites in the pEV-vrf plasmids. The HTLV-III sequences (600 bp) between the *EcoRI* and the *KpnI* site were then removed from the intermediate construction as shown in Figure 3. These plasmid constructions were carried out with all three pEV-vrf plasmids so that subsequent deletions could be made and the correct reading frame maintained. In addition, the constructions made in the incorrect reading frames served as important controls in the expression experiments described below.

In another embodiment of this invention, a second set of expression plasmids were constructed in a similar fashion by deleting sequences between *EcoRI* and *StuI* sites which occur 483 bp downstream of the env gene. Again these deletions (designated pEV/env 205-640) were made in all three reading frames. The translation termination codon used in all of the env expression plasmids is presumably an in-frame TAA located 23 bp downstream of the *HindIII* site in the plasmid. Thus, 8 amino acid residues at the carboxyl terminus are encoded by pBR322 sequences contained within the pEV-vrf expression plasmids.

Expression of ENV AIDS

There are several approaches to screen for bacterial clones containing env AIDS cDNA. These include, for example, RNA selection hybridization, differential hybridization, hybridization with a synthetic probe and screening for clones that produce the desired protein by immunological or biological assays. Two methods are available to screen using immunological assay: screening of bacterial colonies for the presence of protein using antibody; and, preferably, the bacterial lysates are electrophoresed, blotted onto a nitrocellulose paper and then probed with the antibody.

In a preferred embodiment of this invention, cultures of the *E. coli* strain MC 1061 transformed with pRK248cits and the pEV 1, 2, or 3/env 44-640 (or pEV 1, 2 or 3/env 205-640) were grown in M9 medium at 30° C to mid-log phase and then induced by shifting to 42° C for 2 hr. Samples of the bacterial cultures were then taken and subjected to SDS-polyacrylamide gel electrophoresis, followed by Western blot analysis to detect env proteins. The protein blots were treated with antisera to env AIDS proteins isolated either from immunized rabbits or from AIDS patients previously shown to contain high titer antibodies to AIDS antigens. This was followed by incubation with ¹²⁵I-labelled Staphylococcus aureus protein A, washing and autoradiography. Similar results were obtained with both sera except that the human serum was found to contain much higher titers of anti-HTLV-III antibodies and was devoid of all background reactivity with the *E. coli* proteins. For this reason human antibodies were used in all subsequent characterization.

Figure 4 shows the pattern of reactivity of the env AIDS proteins synthesized in bacteria (recombinant proteins) with anti-HTLV-III antibodies. The open reading frame in pEV3/env 44-640 encodes a protein that should migrate as a 68 Kd band on the gel. In fact, a 68 Kd band is observed in the lane corresponding to the induced cells containing pEV3/env 44-640 (lane C). However, in addition to the 68 Kd band, these cells synthesized proteins of 35 Kd, 25 Kd and 17 Kd which specifically cross-reacted with anti-HTLV-III antibodies. No HTLV-III cross-reacting bands are evident in the uninduced control (Lane b) or in a second negative control sample (Lane a) of induced cells containing a plasmid that directs the synthesis of v-bas p21 oncogene product [Lacal, J.C. et al., supra]. The appearance of multiple bands synthesized from the env gene sequences was an unexpected result. Another unexpected result was the synthesis of env gene products from the plasmid (pEV1/env 44-640) where the insert was placed in the wrong reading frame with respect to the initiator codon immediately downstream of the P_L promoter (Lane d). In this case, *E. coli* cells containing plasmid pEV1/env. 44-640 synthesized a 63 Kd protein in addition to the 35 Kd, 25 Kd and 17 Kd proteins. These results could be readily explained when the nucleotide sequence of the envelope gene (Fig. 1) was examined. About 155

bases downstream to the KpnI site is an ATG codon which appeared to be utilized for the synthesis of the env gene product by the expression plasmid pEV1/env 44-640. Internal translation initiation is also the likely explanation for the appearance of the 35Kd, 25Kd and 17Kd proteins. Initiation codons which are preceded by so-called Shine-Dalgarno sequences (AGGA) are found within the env coding region at locations that are consistent with the sites of the observed protein products.

To confirm the above interpretation and to rule out the possibility that the smaller proteins are not formed as a result of premature termination or from proteolytic cleavage of the larger product, another deletion mutant in which sequences between the KpnI and StuI sites were deleted were constructed. This expression plasmid contains the coding sequences from amino acid positions 205-640 which could code for a protein of 49 Kd. Analysis of the proteins induced from *E. coli* harboring this plasmid verified that, in fact, these cells synthesize a 49 Kd protein in addition to the 35 Kd, 25 Kd and 17 Kd proteins (lane e, Fig. 4). From these results, it was concluded that pEV3/env 44-640 expression plasmid directs the synthesis of a 68 Kd protein in addition to several additional smaller polypeptides (i.e., 35Kd, 25Kd and 17Kd) produced from all of the env expression plasmids resulting from internal translation initiation within the env gene.

Screening of AIDS SERA

Because anti-HTLV-III antibodies are found in more than 90% of the AIDS patients, it was of interest to see if the bacterially synthesized env gene products could be used as diagnostic tools for the detection of these antibodies. For this analysis, total cell protein from an induced bacterial culture was fractionated by SDS-PAGE and transferred to a nitrocellulose filter by Western blotting technique. Strips of the filter containing transferred proteins were reacted with 1000-fold diluted human sera, and the antigen-antibody complexes formed were detected by incubation of the strips with 125-I-labelled *Staphylococcus aureus* protein A followed by autoradiography. Prominent bands corresponding to reaction of the antibody to the 68 Kd, 35 Kd, 25 Kd and 17 Kd proteins were consistently observed when the serum used was from patients with AIDS syndrome. The results of such assays with different human sera are presented in Figure 5. The negative controls used were normal human sera and serum from a patient with HTLV-I infection. No reaction was observed with sera from healthy individuals or from HTLV-I infected individuals. The patient sera were derived from all parts of the United States including California and all AIDS patients' sera tested so far were found to be positive. The results suggest that these antibodies are mainly directed against the protein back-bone of the molecule.

It appears, therefore, that the env gene products constitute the best diagnostic reagents for the detection of AIDS associated antibodies. The env gene product of the instant invention encompasses a large portion of the protein molecule and contains both the conserved and divergent portions of the molecule. In spite of the divergence observed between HTLVIII and ARV-2 sequences the recombinant env proteins of the instant invention synthesized by the bacteria react with AIDS patient sera derived from both geographical locations of the United States. One hundred percent (100%) of AIDS patient sera (50 individual samples, 25 derived from the East Coast of the United States and 25 derived from California) tested showed high reactivity. This is strong evidence for the presence of conserved epitopes within the molecule against which the immune system could mount an antibody reaction. The human immune system may thus be mounting an immune response against conserved epitopes of the envelope molecule, as suggested by the reactivity of the AIDS patient sera. The observed divergence between various isolates of HTLV-III thus may not pose a problem for the use of recombinant protein as a vaccine. The 68Kd protein is ideally suited for such a purpose since it encompasses a large portion of the gene product and has the unique structural feature of containing both the extracellular hydrophilic region and the membrane associated hydrophobic regions. This structural feature makes it well suited for encapsulation into liposomes which have been used as vehicles for vaccination against other vital envelope proteins.

Based on these discoveries it is proposed that in the practice of screening blood for AIDS only AIDS envelope protein or a variant of said protein be utilized. Utilizing the env AIDS protein of the instant invention, human blood can be screened for the presence of antibodies to the AIDS virus. This and other techniques are readily determined, once, as taught for the first time by the present invention, the envelope AIDS protein has been recognized to be the envelope protein of the etiologic agent of AIDS. The foregoing and other objects, features and advantages of the invention will be apparent from the following examples of preferred embodiments of the invention.

Example 1

Molecular cloning and nucleotide sequence analysis of the HTLV-III proviral genome.

The integrated proviral genome of HTLV-III was recently cloned from the genomic DNA of H9 cells infected with HTLV-III [Shaw, G.M. et al., supra]. The proviral genome which was obtained by using XbaI digested H9/HTLV-III DNA contained two internal EcoRI sites within the viral genome and two additional sites in the cloning vector λ JI. These sites were used for further subcloning of the three DNA fragments of 5.5Kb, 4.5Kb and 1.1Kb into pBR322 (ATCC No. 37017). Nucleotide sequence analysis of the proviral genome was determined by the chemical degradation method of Maxam, A.M. and Gilbert, W., "Sequencing end-labelled DNA with base-specific chemical cleavages", Meth. Enzymol.

65, 499-560 (1980). For the sequence analysis, DNA inserts from the three subclones were isolated by electroelution and further cleaved with appropriate restriction enzymes. The DNA fragments were labelled at their 5' ends with γ -³²P-ATP using polynucleotide kinase, or at their 3' ends with α -³²P-NTP by filling in with DNA polymerase I (Klenow fragment). The DNA fragments labelled at the two ends were cleaved with a second enzyme and the fragments labelled at a single end were purified on 5% acrylamide gels and used for sequence analysis. For the sequence analysis of the env gene, a shotgun approach was utilized where the 4.5 EcoRI fragment was cleaved with one of the following enzymes: BglII, HindIII, XhoI, AvalI, HinfI and Sau3A and the restriction fragments labeled and sequenced as described above. The nucleotide sequence of the envelope gene used in the present invention is shown in Figure 1.

10 Example 2

Construction of pEV/env 44-640

pRC2 is a derivative of pBR322 containing a unique Bgl II site adjacent (on the amp^R side) to the EcoRI site in the plasmid. This plasmid was constructed in the following manner. 20 μ g of pBR322 plasmid DNA were digested with EcoRI and then split into two reactions. In one, the protruding 5' single-stranded termini were removed with S1 nuclease; in the other reaction, the termini were filled-in by incorporating deoxynucleotides with the Klenow fragment of DNA polymerase I. Both reactions were terminated by phenol extraction followed by ethanol precipitation. Approximately 1 μ g of DNA from each reaction was mixed with 90 pmoles of phosphorylated BglII linkers (CAGATCTG, purchased from Collaborative Research) and incubated with T4 DNA ligase at 15° C for 18 hours. The ligation products were then digested with BglII and PstI and subjected to gel electrophoresis in 1% agarose. The 3600 bp and 760 bp fragments from both reactions were recovered from the gel. For the construction of pRC2, the 3600 bp from the Klenow reaction was ligated to the 760 bp fragment from the S1 reaction. To construct a plasmid with the BglII site on the other side of EcoRI (tet^R side), designated pRC1, the 3600 bp fragment from the S1 reaction was ligated to the 760 bp fragment from the Klenow reaction. E. coli strain RRI (ATCC No. 31343) was transformed with the ligation mixtures, and transformants were selected on LB agar plates containing 50 μ g/ml ampicillin. Transformants containing the expected plasmid constructions were identified by restriction analysis of the isolated plasmid DNA. DNA sequence analysis confirmed that the S1 nuclease treatment precisely removed the 5' single-stranded termini.

pRC23 (see Figure 7) was constructed by inserting into pRC2 a 250 bp BglII-HaeIII fragment containing the λ P_L promoter joined to a pair of complementary synthetic oligonucleotides comprising a model ribosome-binding site (RBS). The HaeIII site is located within the 5' non-coding region of the λ N gene 115 bp downstream of the P_L transcriptional initiation site. Approximately 1 μ g of a 450 bp BglII-HpaI fragment isolated from phage λ DNA was digested with HaeIII. 200 ng of the resulting digestion products were mixed with 60 pmoles each of phosphorylated synthetic oligonucleotides containing the model RBS. The ligated molecules were digested with BglII and EcoRI and separated on a 5% polyacrylamide gel. The 270 bp ligation product was recovered from the gel, mixed with gel purified pRC2 vector that had been digested with BglII and EcoRI, and incubated with T4 DNA ligase at 15° C for 15 hours. The ligation mixture was used to transform strain RRI(pRK248CIts). Transformants selected on ampicillin-containing medium were screened by restriction analysis of the isolated plasmid DNA. The expected plasmid construction, pRC23, was confirmed by further restriction enzyme digestions and by DNA sequence analysis across the EcoRI junction (Fig. 7).

For the construction of the pEV-vrf set of plasmids (see Figure 8), plasmid pRC23 was digested with EcoRI and HindIII and the pRC23/EcoRI-HindIII vector isolated by preparative agarose gel electrophoresis. The mixture of synthetic oligonucleotides (32, 33, and 34 nucleotides) was combined with the mixture of the complementary sequences, heated to 58° C for 5 minutes in 150 mM NaCl, and cooled slowly to allow annealing. 0.1 pmoles of the synthetic duplexes were added to 0.07 pmoles of the pRC23/EcoRI-HindIII vector and incubated with T4 DNA ligase at 15° C for 15 hours. Strain RRI (λ cl857) was transformed with the ligation products. Six ampicillin resistant transformants were selected for DNA sequence analysis. Of the six, two contained the expected sequence for pEV-vrf1, one for pEV-vrf2, and three for pEV-vrf3 (Fig. 3).

For the expression of the AIDS env gene, one μ g of a 2400 bp EcoRI - HindIII DNA fragment, which was isolated from the cloned HTLV-III proviral genome by preparative agarose gel electrophoresis, was mixed with 0.1 μ g of EcoRI - HindIII digested vector DNA (pEV-vrf1, -2, or -3). After heating at 65° C for 3 minutes, the mixtures were chilled on ice, and 20 μ l ligation reactions were assembled, containing 50 mM Tris-HCl (pH 7.4), 10 mM MgCl₂, 10 mM DTT, 0.3 mM ATP, and 200 units of T₄ DNA ligase. After incubation at 15° C for 4 hours, the reactions were terminated by heating at 65° C for 5 minutes. The ligation products were used to transform E. coli strain MC1061 containing plasmid pRK248CIts. Transformants were selected on Luria broth agar containing 50 μ g/ml ampicillin at 30° C for 18 hours. Plasmid DNA was isolated from 1 ml of each culture and subjected to restriction analysis. All 12 isolates contained the expected plasmid construction. These intermediate constructions were then used to make pEV1, -2, and -3/env 44-640 by deleting the 600 bp between the EcoRI and KpnI sites as described below.

Approximately 0.5 μ g of plasmid DNA was digested with KpnI and EcoRI. The resulting termini were then treated with the Klenow fragment of DNA polymerase I in the presence of all four deoxyribonucleotides (at 100 μ M) at 37° C for

30 minutes. This step results in the "filling-in" of the 5' overhang of the EcoRI terminus and the removal of the 3' overhang of the KpnI terminus. Upon recircularization of the linear plasmid and blunt-end ligation of these termini, an EcoRI site is regenerated. Transformants containing plasmids with the expected deletion were identified by restriction analysis.

- 5 A second set of deletion derivatives, designated pEV/env 205-640 was constructed in a similar fashion. A portion of the linear plasmid that had been digested with EcoRI and KpnI and treated with Klenow, as described above, was further digested with StuI. Again, upon recircularization and blunt-end ligation, the EcoRI site was regenerated; however, an additional 483 bp of env coding sequences were removed.

10 Example 3

Bacterial Growth and Induction of env Gene Expression

- 15 Cultures of E. coli strain MC 1061 transformed with plasmid pRK248cIts and the pEV1, -2, or -3/env plasmids were grown in M9 medium containing 0.5% glucose and 0.5% casamino acids at 30° C to mid-log phase and then induced by shifting to 42° C for 2 hr. The cells were collected by centrifugation and processed as described in Examples 4 and 5.

Example 4

20 Expression and Purification of Env AIDS

A homogeneous recombinant viral env AIDS was purified according to the following procedure. The env AIDS protein expressed by a microbe tends to associate with the membrane fractions of the host microbe, principally the inner membrane of the microbe. The following purification method was designed to deal with this finding.

- 25 This purification method comprises:

- (a) lysis of transformed microbial cells producing recombinant env AIDS protein;
- (b) separation of env AIDS associated cellular membranes from other cellular components;
- (c) extraction of env AIDS from associated membranes; and
- 30 (d) chromatographic purification of the resultant extraction solution containing env AIDS to yield a substantially pure recombinant viral env protein.

More specifically, the preferred purification method for the preparation of substantially pure recombinant viral env protein comprises:

- 35 (a) cultivating a transformed organism containing a DNA sequence which codes for viral env protein;
- (b) causing a culture of the transformed organism of step (a) to accumulate the env protein;
- (c) lysing the culture of transformed organisms of step (b) to form a cell lysate mixture;
- (d) isolating the cell membrane components of the cell lysate mixture of step (c);
- 40 (e) washing the isolated cell membrane components with an extraction solution to yield a wash solution containing env protein; and
- (f) chromatographically purifying the wash solution of step (e) to yield a substantially pure env AIDS protein.

- 45 In carrying out this method it is preferred that the cells be lysed by sonification, although it is foreseeable that other known methods such as enzyme or mechanical lysis could also be used. It is preferred that the cell membrane component, specifically the inner and outer membranes, be isolated from other cellular components by methods such as centrifugation. It has been found that env AIDS expressed by the transformed microorganism tends to become associated with the cellular membranes. Therefore, isolation of these membranes during the purification process ensures high purification levels and high purity env AIDS at the end of the purification procedure.

- 50 Once the cell membranes are isolated from the lysate mixture, they are washed with an extraction solution, preferably salt solutions and a detergent to yield a second solution containing approximately 50% env AIDS protein. Preferably the cell membranes are washed in four separate steps with the salt solutions and detergent although it is foreseeable that certain of these steps could be combined, rearranged or eliminated. The first step of washing the cell membrane may be done with a salt solution, preferably 1M NaCl. In the second step the cell membrane is washed with a detergent solution, preferably 1% Triton X-100. In the third step, the cell membrane is washed with another salt solution, 1.75M to 3.5M guanidine HCl. The final wash is also with a salt solution preferably about 7M Guanidine HCl. The wash solution which results from the fourth and final wash comprises about 50% env AIDS.

The final 50% env AIDS wash solution is then further purified by a chromatography step, preferably reverse phase high performance liquid chromatography (HPLC). The HPLC step yields env AIDS protein in a substantially 100% pure

form. It is also foreseeable that monoclonal antibody affinity chromatography columns utilizing env AIDS polyclonal or monoclonal antibodies, could be used as an alternative to HPLC.

Example 5

Polyacrylamide gel electrophoresis and Western blot analysis

Cells were lysed by resuspending the cell pellets (approximately 10^8 cells) in TG buffer (10 mM Tris, pH 7.4, 10% glycerol), mixed with an equal volume of 2 x sample buffer [Laemmli, U.K., "Cleavage of Structural Proteins During the Assembly of the Head of Bacteriophage T4", Nature 227, 680-685 (1970)] and incubated at 95° C for five (5) minutes. Cell debris were pelleted by centrifugation and the cleared lysates were subjected to SDS-PAGE analysis [Laemmli, U.K., supra]. For Western blot analysis, the proteins from the acrylamide gel were electroblotted onto a 0.1 µm nitrocellulose membrane (Schleicher and Schuell) for 16 hr at 50V, in 12.5 mM Tris, 96 mM glycine, 20% methanol, 0.01% SDS at pH 7.5. Processing of the blot was carried out using the methods described by Towbin, H. et al. ["Electrophoretic Transfer of Proteins From Polyacrylamide Gels to Nitrocellulose Sheets: Procedure and Some Applications", Proc. Natl. Acad. Sci. U.S.A., 76, 4350-4354, (1979)]. For treatment with the human sera, the blots were incubated with a 1000 fold dilution of the sera in antibody buffer (20 mM sodium phosphate buffer, pH 7.5, containing 0.5 M NaCl, 1% BSA and 0.05% Tween 20) for 2-6 hr. The blots were then washed twice with phosphate buffered saline containing 0.05% Tween 20 and then incubated with 125-I-labelled Staphylococcus aureus protein A for an additional period of 1 hr. The blot was then washed twice in PBS-Tween 20 buffer, dried and autoradiographed.

Example 6

Immunization with Env Protein of AIDS Virus

It is clear that in spite of the divergence observed between HTLVIII and ARV-2 sequences, the recombinant proteins synthesized by the bacteria react well with AIDS patients' sera derived from both geographical locations of the United States. One hundred percent (100%) of the AIDS patients' sera tested showed high reactivity (50 individual samples, 25 from the east coast of the United States and 25 from the west coast of the United States). Thus all the env proteins contain at least one conserved epitope. All of the human sera from AIDS patients tested contained antibodies to the env proteins of the instant invention. This strongly suggests that these env proteins with the conserved epitopes would be immunogenic in man.

It will be readily appreciated that the env proteins of the instant invention can be incorporated into vaccines capable of inducing protective immunity against the AIDS virus. By methods known in the art, the specific amino acids comprising the epitopes of the env protein may be determined. Peptides may then be synthesized, comprising an amino acid sequence corresponding to an epitope of an env AIDS protein either in monomeric or multimeric form. These synthetic peptides may then be incorporated into vaccines capable of inducing protective immunity against AIDS virus. Techniques for enhancing the antigenicity of such peptides include incorporation into a multimeric structure, binding to a highly immunogenic protein carrier, for example, keyhole limpet hemocyanin, or diphtheria toxoid, and administration in combination with adjuvants or any other enhancers of immune response. In addition, the vaccine composition may comprise antigens to provide immunity against other diseases in addition to AIDS.

An amino acid sequence corresponding to an epitope of an env protein either in monomeric or multimeric form (peptide) may be obtained by chemical synthetic means or by purification from biological sources including genetically modified microorganisms or their culture media. The peptide may be combined in an amino acid sequence with other peptides including fragments of other proteins, as for example, when synthesized as a fusion protein, or linked to other antigenic or non-antigenic peptides of synthetic or biological origin. The term "corresponding to an epitope of an env protein" will be understood to include the practical possibility that, in some instances, amino acid sequence variations of a naturally occurring peptide may be antigenic and confer protective immunity against AIDS infection. Possible sequence variations include, without limitation, amino acid substitutions, extensions, deletions, interpolations and combinations thereof. Such variations fall within the contemplated scope of the invention provided the peptide containing them is antigenic and antibodies elicited by such peptide cross-react with naturally occurring env protein or non-variant repeated peptides of env protein, to an extent sufficient to provide protective immunity when administered as a vaccine. Such vaccine compositions will be combined with a physiologically acceptable medium. The size and shape of epitopes found in carbohydrate antigens have been extensively studied, but less is known about the structure of epitopes from protein molecules. Some epitopes of protein antigens have been defined at the level of their tertiary structure. In every instance, the epitopes were formed not by the primary sequences alone, but by the juxtaposition of residues brought together by the folding of the polypeptide chain(s) of the native molecule. In addition, the structure of the 68Kd env protein of the instant invention makes it particularly well suited for use as a vaccine. The 68Kd env protein comprises a large portion of the gene product which (a) was shown to be reactive with all the AIDS sera tested; and (b) has the

unique structural feature of containing both an extracellular hydrophilic region and the transmembrane hydrophobic regions. The latter structural feature makes it well suited for use as a vaccine using liposome encapsulation to create a vehicle for administration.

Routes of administration, antigen dose, number and frequency of injections are all matters of optimization within the scope of ordinary skill in the art, particularly in view of the fact that there is experience in the art in providing protective immunity by the injection of other related antigens to provide immunity in other viral infections. It is anticipated that the principal value of providing immunity to AIDS infection will be for those individuals who have had no previous exposure to AIDS, e.g., individuals who are in the high risk population, such as homosexuals, drug addicts and people from Haiti and Central America and individuals who may be receiving blood transfusions. It is also anticipated that temporary immunity for infants may be provided by immunization of mothers during pregnancy.

Example 7

Diagnostic Test for AIDS

It is clear that the env gene proteins of the instant invention may be used as diagnostic reagents for the detection of AIDS-associated antibodies. It is also apparent to one of ordinary skill that a diagnostic assay for AIDS using polyclonal or monoclonal antibodies to the AIDS env protein of the instant invention may be used to detect the presence of the AIDS virus in human blood. In one embodiment a competition immunoassay is used where the antigenic substance, in this case the AIDS virus, in a blood sample competes with a known quantity of labelled antigen, in this case labelled AIDS env protein, for a limited quantity of antibody binding sites. Thus, the amount of labelled antigen bound to the antibody is inversely proportional to the amount of antigen in the sample. In another embodiment, an immunometric assay may be used wherein a labelled AIDS-env antibody is used. In such an assay, the amount of labelled antibody which complexes with the antigen-bound antibody is directly proportional to the amount of antigen (AIDS virus) in the blood sample. In a simple yes/no assay to determine whether the AIDS virus is present in blood, the solid support is tested to detect the presence of labelled antibody. In another embodiment, monoclonal antibodies to AIDS env protein may be used in an immunometric assay. Such monoclonal antibodies may be obtained by methods well known in the art, particularly the process of Milstein and Kohler reported in Nature 256, 495-497 (1975).

The immunometric assay method is as follows: Duplicate samples are run in which 100 μ l of a suspension of antibody immobilized on agarose particles is mixed with 100 μ l of serum and 100 μ l of soluble 125 I-labelled antibody. This mixture is for specified times ranging from one quarter hour to twenty four hours. Following the incubation periods the agarose particles are washed by addition of buffer and then centrifuged. After removal of the washing liquid by aspiration, the resulting pellet of agarose particles is then counted for bound 125 I-labelled antibody. The counts obtained for each of the complexes can then be compared to controls.

While the invention has been described in terms of certain preferred embodiments, modifications obvious to one with ordinary skill in the art may be made without departing from the scope of the invention. For example, it is understood that the env AIDS DNAs described herein represent only the precise structure of two naturally occurring gene segments. It is expected that slightly modified alleles will be found encoding for similarly functioning proteins, and such gene segments and proteins are considered to be equivalents for the purpose of this invention. It is also suspected that other variants in addition to those described herein will be found and that the envelope protein of said variants will differ slightly. These variant envelope proteins are likewise considered within the scope of the invention. DNA having equivalent codons is considered within the scope of the invention, as are synthetic gene segments that encode homologous proteins of the viral envelope.

Various features of the invention are set forth in the following claims.

Claims

Claims for the following Contracting States : BE, CH, DE, FR, GB, IT, LI, NL, SE

1. An envelope protein fragment of an acquired immune deficiency syndrome (AIDS) virus, essentially free of other proteins, with the amino acid sequence:

ValTrpLysGluAla
 ThrThrThrLeuPheCysAlaSerAspAlaLysAlaTyrAspThrGluValHisAsnValTrpAlaThr
 HisAlaCysValProThrAspProAsnProGlnGluValValLeuValAsnValThrGluAsnPheAsn
 5 METTrpLysAsnAspMETValGluGlnMETHisGluAspIleIleSerLeuTrpAspGlnSerLeuLys
 ProCysValLysLeuThrProLeuCysValSerLeuLysCysThrAspLeuLysAsnAspThrAsnThr
 AsnSerSerSerGlyArgMETIleMETGluLysGlyGluIleLysAsnCysSerPheAsnIleSerThr
 SerIleArgGlyLysValGlnLysGluTyrAlaPhePheTyrLysLeuAspIleIleProIleAspAsn
 10 AspThrThrSerTyrThrLeuThrSerCysAsnThrSerValIleThrGlnAlaCysProLysValSer
 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 15 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 20 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 25 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 30 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

CysProLysValSer
 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 40 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 45 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 50 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 55 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

METArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

METTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

METArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer.

2. An expression vector comprising a gene coding for an envelope protein fragment of an AIDS virus as defined in claim 1 downstream of a promoter sequence enabling transcription, translation and thus expression of said envelope protein fragment in a host cell.
3. An expression vector according to claim 2, wherein said gene coding for an envelope protein fragment of an AIDS virus is a gene comprising the nucleotide sequence:

GTGTGGAAGGAAGCA
 ACCACCACTCTATTTTGTGCATCAGATGCTAAAGCATATGATACAGAGGTACATAATGTTTGGGCCACA
 CATGCCTGTGTACCCACAGACCCCAACCCACAAGAAGTAGTATTGGTAAATGTGACAGAAAATTTTAAC
 5 ATGTGGAAAAATGCATGGTAGAACAGATGCATGAGGATATAATCAGTTTATGGGATCAAAGCCTAAAG
 CCATGTGTAAAAATTAACCCCACTCTGTGTTAGTTTAAAGTGCACTGATTGAAGAAATGATACTAATACC
 AATAGTAGTAGCGGGAGAAATGATAATGGAGAAAGGAGAGATAAAAACTGCTCTTTCAATATCAGCACA
 AGCATAAGAGGTAAGGTGCAGAAAGAATATGCATTTTTTTTATAAACTGATATAATACCAATAGATAAT
 10 GATACTACCAGCTATACGTTGACAAGTTGTAACACCTCAGTCATTACACAGGCCTGTCCAAAGGTATCC
 TTTGAGCCAAATTCCTATACATTATTGTGCCCCGGCTGGTTTTGCGATTCTAAAAATGTAATAATAAGACG
 TTCAATGGAAACAGGACCAATGACAAATGTGACACAGTACAAATGTACACATGGAAATTAGGCCAGTAGTA
 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAAGAGGTAGTAATTAGATCTGTCAATTTTCAG
 GACAAATGCTAAAAACCATAAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCAAACAAC
 15 AATACAAGAAAAAAATCCGTATCCAGAGGGGACCAGGGAGAGCATTGTTACAATAGGAAAAATAGGA
 AATATGAGACAAGCACAATTGTAACATTAGTAGAGCAAAATGGAAATGCCACTTTAAAAACAGATAGCTAGC
 AAATTAAGAGAACAAATTTGAAATAATAAAACAATAATCTTTAAGCAATCTTCAGGAGGGGACCCAGAA
 ATTGTAAOCACAGTTTTAATTGTGGAGGGGAATTTTTCTACTGTAATTCACACAATGTTTTAATAGT
 ACTTGGTTTTAATAGTACTTGGAGTACTGAAGGGTCAAAATACACTGAAGGAAGTGACACAATCACACTC
 20 CCATGCAGAAATAAAACAATTTATAAACAATGTCAGGAAGTAGGAAAAGCAATGTATGCCCCCTCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGACGGACAATTGGAGAAGTGAATTAATAAAA
 TATAAAGTAGTAAAAATGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGACAGAGA
 25 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACCGTGACGGTACAGGCCAGACAATTAATTGTCTGGTATAGTCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGATTTGGGGTTGCTCTGGA
 30 AAATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGGA
 ATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

or an equivalent thereof, coding for said envelope protein fragment.

- 35 4. An expression vector according to claim 2, wherein said gene coding for an envelope protein fragment of an AIDS virus is a gene comprising the nucleotide sequence:

40

45

50

55

TGTCCAAAGGTATCC

TTTGAGCCAATTCCCATACATTATTGTGCCCCGGCTGGTTTTGCGATTCTAAAAATGTAATAATAAGACG
 TTCAATGGAACAGGACCATGTACAAATGTCAGCACAGTACAATGTACACATGGAATTAGGCCAGTAGTA
 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAAGAGGTAGTAATTAGATCTGTCAATTTACAG
 GACAATGCTAAAAACCATAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCAACAAC
 AATACAAGAAAAAAATCCGTATCCAGAGGGGACCAGGGAGAGCATTGTTTACAATAGGAAAAATAGGA
 AATATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAAAACAGATAGCTAGC
 AAATTAAGAGAACAAATTTGGAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACGCACAGTTTTAATTGTGGAGGGGAATTTTCTACTGTAATTCAACACAACCTGTTTAAATAGT
 ACTTGGTTTTAATAGTACTTGGAGTACTGAAGGGTCAAAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCAGAATAAAACAATTTATAAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAAATCTCTGGAACAGATTTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

or an equivalent thereof, coding for said fragment.

5. An expression vector according to claim 2, wherein said gene coding for an envelope protein fragment of an AIDS virus is a gene comprising the nucleotide sequence:

ATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAAAACAGATAGCTAGC
 AAATTAAGAGAACAAATTTGGAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACGCACAGTTTTAATTGTGGAGGGGAATTTTCTACTGTAATTCAACACAACCTGTTTAAATAGT
 ACTTGGTTTTAATAGTACTTGGAGTACTGAAGGGTCAAAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCAGAATAAAACAATTTATAAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAAATCTCTGGAACAGATTTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

or an equivalent thereof, coding for said envelope protein fragment.

6. An expression vector according to claim 2, wherein said gene coding for an envelope protein fragment of an AIDS virus is a gene comprising the nucleotide sequence:

ATGTATGCCCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGA CAATTGGAGAAGTGAATTATATAAA
 5 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACCGTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAT
 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 10 GCAAGAACTCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

15 or an equivalent thereof coding for said envelope protein fragment.

7. An expression vector according to claim 2, wherein said gene coding for an envelope protein fragment of an AIDS virus is a gene comprising the nucleotide sequence:

ATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACCGTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAT
 25 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAACTCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

30

8. An expression vector according to any one of claims 2 to 7, which is a plasmid capable of replication in gram-negative and/or gram-positive bacteria.

35

9. An expression vector according to claim 8 which is capable of replication in an E. coli strain.
 10. An expression vector according to claim 8 which is capable of replication in a B. subtilis strain.

40

11. The expression vector pEV1. -2. or -3/env 44-640.

12. The expression vector pEV1. -2. or -3/env 205-640.

13. A transformant carrying an expression vector as claimed in any one of claims 2 to 12.

45

14. A transformant according to claim 13 which is an E. coli strain.

15. A transformant according to claim 14 which is an E. coli MC 1061 strain.

50

16. A transformant according to claim 13 which is a B. subtilis strain.

17. A transformant according to claim 13 which is a mammalian cell.

55

18. A method of producing an envelope protein fragment of an acquired immune deficiency syndrome virus as claimed in claim 1 comprising:

transforming a host cell with an expression vector as claimed in any one of claims 2 to 12; culturing said host cell so that said AIDS env protein fragment is expressed; and
 extracting and isolating said AIDS env protein fragment.

19. A method according to claim 18, wherein the expression vector is pEV1, -2 or -3/env 44-640.

20. A method according to claim 18, wherein the expression vector is pEV1, -2 or -3/env 205-640.

5 21. A method of testing human blood for the presence of antibodies to the viral etiologic agent of AIDS which comprises mixing a composition containing an envelope protein fragment of an AIDS virus as claimed in claim 1 with a sample of human blood and determining whether said envelope AIDS protein fragment binds to AIDS antibodies present in the blood sample.

10 22. A method according to claim 21 which comprises the use of the Western Blotting Analysis.

23. A method according to claim 21 which comprises the use of an ELISA-technique, wherein an envelope protein fragment of an AIDS virus as claimed in claim 1 is coated on a solid phase and contacted with the sample and after washing contacted with an enzyme-labeled non-human IgG.

15 24. A method according to claim 21, wherein the Double-Antigen-Method is used.

25. A method for the determination of AIDS virus, wherein antibodies against an envelope protein fragment of an AIDS virus according to claim 1 are used.

20 26. A method according to claim 25, wherein the antigen in the sample and a protein fragment as claimed in claim 1 in labeled form compete with an antibody against a protein fragment as claimed in claim 1.

25 27. A method according to claim 25, wherein a sandwich method is performed using two antibodies against a protein fragment as claimed in claim 1.

28. A method according to claim 27, wherein one antibody is on a solid phase and the other antibody is labeled.

29. A method according to claim 27, wherein two different monoclonal antibodies are used.

30 30. A vaccine eliciting immunity to AIDS comprising as an active ingredient a protein fragment as claimed in claim 1.

31. Antibodies raised against a protein fragment as claimed in claim 1.

35 32. The antibodies of claim 31 which are monoclonal antibodies.

33. The use of a protein fragment as claimed in claim 1 for the preparation of a protective immunisation vaccine.

34. The use of a protein fragment as claimed in claim 1 for testing human blood for the presence of AIDS virus.

40 **Claims for the following Contracting State : AT**

1. A process for the preparation of an envelope protein fragment of an acquired immune deficiency syndrome (AIDS) virus, essentially free of other proteins, comprising:

45 transforming a host cell with an expression vector comprising a gene coding for an envelope protein fragment of an AIDS virus with the amino acid sequence:

50

55

ValTrpLysGluAla
 ThrThrThrLeuPheCysAlaSerAspAlaLysAlaTyrAspThrGluValHisAsnValTrpAlaThr
 HisAlaCysValProThrAspProAsnProGlnGluValValLeuValAsnValThrGluAsnPheAsn
 5 METTrpLysAsnAspMETValGluGlnMETHisGluAspIleIleSerLeuTrpAspGlnSerLeuLys
 ProCysValLysLeuThrProLeuCysValSerLeuLysCysThrAspLeuLysAsnAspThrAsnThr
 AsnSerSerSerGlyArgMETIleMETGluLysGlyGluIleLysAsnCysSerPheAsnIleSerThr
 SerIleArgGlyLysValGlnLysGluTyrAlaPhePheTyrLysLeuAspIleIleProIleAspAsn
 10 AspThrThrSerTyrThrLeuThrSerCysAsnThrSerValIleThrGlnAlaCysProLysValSer
 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 15 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 20 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 25 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 30 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

35
 CysProLysValSer
 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 40 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 45 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 50 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 55 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

METArgGlnAlaHisCysAsnIle SerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 5 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 10 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 15 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

METTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 25 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 30 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

or

METArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 40 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

downstream of a promoter sequence enabling transcription, translation and expression of said envelope pro-
 tein fragment in said host cell; culturing said host cell so that said envelope protein fragment of an AIDS virus
 is expressed; and extracting and isolating said envelope protein fragment of an AIDS virus.

2. A process according to claim 1, wherein the host cell is a bacterium.
3. A process according to claim 2, wherein the bacterium is *E. coli*.
4. A process according to claim 3, wherein the plasmid is pEV1, -2, or -3/env 44-640.
5. A process according to claim 3, wherein the plasmid is pEV1, -2, or -3/env 205-640.
6. A process for the preparation of an expression vector comprising a gene coding for an envelope protein fragment of an AIDS virus, which process comprises constructing an expression vector having an insertion site, wherein a

gene coding for an envelope protein fragment of an AIDS virus as defined in claim 1 may be inserted which insertion site is downstream of a promoter sequence enabling transcription, translation and thus expression of said envelope protein fragment in a host cell.

- 5 7. A process according to claim 6, characterized in that as said gene coding for an envelope protein fragment of an AIDS virus a gene comprising the nucleotide sequence

GTGTGGAAGGAAGCA

10 ACCACCACTCTATTTTGTGCATCAGATGCTAAAGCATATGATACAGAGGTACATAATGTTTGGGCCACA
CATGCCCTGTGTACCCACAGACCCCAACCCACAAGAAGTAGTATTGGTAAATGTGACAGAAAATTTTAAC
ATGTGGAAAAATGACATGGTAGAACAGATGTCATGAGGATATAATCAGTTTATGGGATCAAAGCCTAAAG
CCATGTGTAAAATTAACCCCACTCTGTGTTAGTTTAAAGTGCCTGATTTGAAGAAATGATACTAATACC
AATAGTAGTAGCGGGAGAATGATAATGGAGAAAGGAGAGATAAAAACTGCTCTTTCAATATCAGCACA
15 AGCATAAGAGGTAAAGGTGCAGAAAGAATATGCATTTTTTATAAACTTGATATAATACCAATAGATAAT
GATACTACCAGCTATACGTTGACAAGTTGTAACACCTCAGTCATTACACAGGCCTGTCCAAAGGTATCC
TTTGAGCCAATTCCCATACATTATGTGCCCCGGCTGGTTTTGCGATTCTAAAATGTAATAATAAGACG
TTCAATGGAACAGGACCATGTACAAATGTCAGCACAGTACAATGTACACATGGAATTAGGCCAGTAGTA
TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAGGTTAGTAATTAGATCTGTCAATTTTCAG
20 GACAAATGCTAAAACCATTAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCCAACAAC
AATACAAGAAAAAAATCCGTATCCAGAGGGGACCAGGGAGAGCATTTGTTACAAATAGGAAAAATAGGA
AATATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAAACAGATAGCTAGC
AAATTAAGAGAACAATTTGGAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
ATTGTAACGCACAGTTTAAATGTTGGAGGGGAATTTTCTACTGTAATTCACACAACCTGTTAATAGT
25 ACTTGGTTTAAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACA
CCATGCAGAATAAAACAATTTATAAAATGTTGGCAGGAAGTAGGAAAAGCAATGTATGCCCCCTCCCATC
AGCGGACAAATTAGATGTTCAATAATATTACAGGGCTGCTATTAAACAAGAGATGGTGGTAATAACAAC
AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
30 GAAAAAAGAGCAGTGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAT
TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
GCAAGAATCCTGGCTGTGAAAGATACCTAAAGGATCAACAGCTCCTGGGATTGGGGTTGCTCTGGA
35 AAATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTTGG
AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

or an equivalent, coding therefore is used.

- 40 8. A process according to claim 6, characterized in that as said gene coding for an envelope protein fragment of an AIDS virus a gene comprising the nucleotide sequence

45

50

55

TGTCCAAAGGTATCC

TTTGAGCCAATTCCCATACATTATTGTGCCCCGGCTGGTTTTCGATTCTAAAAATGTAATAATAAGACG
 TTCAA TCGAACAGGACCATGTACAAATGTCAGCACAGTACAAATGTACACATGGAATTAGGCCAGTAGTA
 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAAGAGGTAGTAATTAGATCTGTCAATTTCAAG
 GACAAATGCTAAAAACCATAATAGTACAGCTGAACACATCTGTAGAAAATTAATTGTACAAGACCCAAACAAC
 AATACAAGAAAAAAATCCGTATCCAGAGGGGACCAGGAGAGCATTGTGTTACAATAGGAAAAATAGGA
 AATATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAAACAGATAGCTAGC
 AAATTAAGAGAACAATTTGGAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACGCACAGTTTAAATTGTGGAGGGGAATTTTCTACTGTAATTCACACAAGTGTAAATAGT
 ACTTGGTTTAAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCAGAAATAAAACAATTTATAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCCCTCCCATC
 AGCGGACAAATTAGATGTTTCAATCAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGCGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACCGTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

or an equivalent coding therefore is used.

9. A process according to claim 6, characterized in that as said gene coding for an envelope protein fragment of an AIDS virus a gene comprising the nucleotide sequence

ATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAAACAGATAGCTAGC
 AAATTAAGAGAACAATTTGGAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACGCACAGTTTAAATTGTGGAGGGGAATTTTCTACTGTAATTCACACAAGTGTAAATAGT
 ACTTGGTTTAAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCAGAAATAAAACAATTTATAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCCCTCCCATC
 AGCGGACAAATTAGATGTTTCAATCAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGCGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACCGTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

or an equivalent coding therefore is used.

10. A process according to claim 6, characterized in that as said gene coding for an envelope protein fragment of an AIDS virus a gene comprising the nucleotide sequence

..
 ATGTATGCCCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCGGAGATCTTCAGACCTGGAGGAGGAGATAGAGGGACAATTGGAGAAGTGAATTATATAAA
 5 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAAACAAT
 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 10 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

15 or an equivalent coding therefore is used.

11. A process according to claim 6, characterized in that as said gene coding for an envelope protein fragment of an AIDS virus a gene comprising the nucleotide sequence

20 ATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAAACAAT
 25 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGCT

30 or an equivalent coding therefore is used.

12. A process according to any one of claims 6 to 11, wherein the expression vector is a plasmid capable of replication in gram-negative bacteria.
- 35 13. A process according to claim 12, wherein the plasmid is capable of replication in an E. coli strain.
14. A process for the preparation of a transformant carrying an expression vector comprising a gene coding for an envelope protein fragment of an AIDS virus, which process comprises transforming a microorganism with an expression vector obtained according to any one of claims 6 to 13 and cultivating the transformed microorganism.
- 40 15. A process according to claim 14, wherein the microorganism is an E. coli strain.
16. A process according to claim 15, wherein the microorganism is an E. coli MC 1061 strain.
- 45 17. A process of testing human blood for the presence of antibodies to the viral etiologic agent of AIDS which process comprises mixing a composition containing an envelope protein fragment of an AIDS virus obtained according to claim 1 with a sample of human blood and determining whether said envelope AIDS protein fragment binds to AIDS antibodies present in the blood sample.
- 50 18. A process according to claim 17 which comprises the use of the Western Blotting Analysis.
19. A process according to claim 17 which comprises the use of an Elisa-technique, wherein an envelope protein fragment of an AIDS virus obtained according to claim 1 is coated on a solid phase and contacted with the sample and after washing contacted with an enzyme-labeled non-human IgG.
- 55 20. A process according to claim 17, wherein the Double-Antigen-Method is used.
21. A process for the determination of AIDS virus, wherein antibodies against an envelope protein fragment of an AIDS virus obtained according to claim 1 are used.

22. A process according to claim 21, wherein the antigen in the sample and a protein fragment obtained according to claim 1 in labeled form compete with an antibody against a protein fragment obtained according to claim 1.
23. A process according to claim 21, wherein a sandwich method is performed using two antibodies against a protein fragment obtained according to claim 1.
24. A method according to claim 23, wherein one antibody is on a solid phase and the other antibody is labeled.
25. A method according to claim 23, wherein two different monoclonal antibodies are used.
26. An envelope protein fragment of an AIDS virus whenever prepared by a process as claimed in any one of claims 1 to 5.
27. An expression vector comprising a gene coding for an envelope protein fragment of an AIDS virus whenever prepared by a process as claimed in any one of claims 6 to 13.
28. A transformant carrying an expression vector comprising a gene coding for an envelope protein fragment of an AIDS virus whenever prepared by a process as claimed in any one of claims 14 to 16.
29. An expression vector comprising a gene coding for an envelope protein fragment of an AIDS virus as defined in claim 1 downstream of a promoter sequence enabling transcription, translation and thus expression of said envelope protein fragment in a host cell.
30. An expression vector according to claim 29, wherein said gene coding for an envelope protein fragment of an AIDS virus is a gene comprising the nucleotide sequence:

GTGTGGAAGGAAGCA

ACCACCACTCTATTTTGTGCATCAGATGCTAAAGCATATGATACAGAGGTACATAATGTTTGGGCCACA
 CATGCCTGTGTACCCACAGACCCCCAACCCACAAGAAGTAGTATTGGTAAATGTGACAGAAAATTTTAAC
 ATGTGGAATAATGACATGGTAGAACAGATGCATGAGGATATAATCAGTTTATGGGATCAAAGCCTAAAG
 CCATGTGTAAATAAACCCTCTGTGTTAGTTTAAAGTGCCTGATTTGAAGAATGATACTAATACC
 AATAGTAGTAGCGGAGAATGATAATGGAGAAAGGAGAGATAAAAACTGCTCTTCAATATCAGCACA
 AGCATAAGAGGTAAGGTGCAGAAAGAATATGCATTTTTTTATAAACTTGATATAATACCAATAGATAAT
 GATACTACCAGCTATACGTTGACAAGTTGTAACACCTCAGTCATTACACAGGCCTGTCCAAAGGTATCC
 TTTGAGCCAATTCCCATACATTATTGTGCCCGGCTGGTTTTTGGGATTCTAAAATGTAATAATAAGACG
 TTCAATGGAACAGGACCATGTACAAAATGTCAGCACAGTACAATGTACACATGGAATTAGGCCAGTAGTA
 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAAGAGGTAGTAATTAGATCTGTCAATTTTCAG
 GACAATGCTAAAACCATAAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCAAACAC
 AATACAAGAAAAAAATCCGTATCCAGAGGGGACCAGGGAGAGCATTGTTTACCAATAGGAAAAATAGGA
 AATATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAAAACAGATAGCTAGC
 AAATTAAGAGAACAATTTGGAATAATAAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACGCACAGTTTTAATTGTGGAGGGGAATTTTTCTACTGTAATTCAACACAACCTGTTTAATAGT
 ACTTGGTTTTAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCAGAAATAAACAAATTTATAACAATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCCCTCCCATC
 AGCGGACAAAATTAGATGTTTCATCAAAATTTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTCGAGGAGGAGATATGAGGGACAAATGCGAGAAGTGAATTAATAAAA
 TATAAAGTAGTAAAAATTGAACCATAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGCTCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTCGGGCATCAAGCAGCTCCAG
 GCAAGAATCTTGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCCTTGAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTG
 AATCACACGACCTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

or an equivalent thereof. coding for said envelope protein fragment.

31. An expression vector according to claim 29, wherein said gene coding for an envelope protein fragment for an AIDS virus is a gene comprising the nucleotide sequence:

TGTCCAAAGGTATCC

5 TTTGAGCCAATTCCCATACATTATTGTGCCCCGGCTGGTTTTGCGATTCTAAAAATGTAATAATAAGACG
 TTCAATGGAACAGGACCATGTACAAATGTCAGCACAGTACAATGTACACATOGAATTAGGCCAGTAGTA
 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAGAGGTAGTAATTAGATCTGTCAATTTCAOG
 GACAAATGCTAAAAACCATAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCCAAC
 10 AATACAAGAAAAAATCCGTATCCAGAOOGGACCAGOGAGAGCATTGTGTTACAAATAGGAAAAATAGGA
 AATATGAGACAAGCCATTGTAACATTAGTAGAGCAAAATOGAATGCCACTTTAAACAGATAGCTAGC
 AAATTAAGAGAACAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACGCACAGTTTTAATTGTGGAGGGGAATTTTCTACTGTAATTCACACAACCTGTTTAATAGT
 ACTTGGTTTAATAGTACTTGGAGTACTGAAOOGTCAAATAACACTGAAOGAAGTGACACAATCACACTC
 15 CCATGCAGAAATAAACCAATTTATAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAAACAAGAGATGGTGTAAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAOOGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTTGAACCAATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 20 GCAGCGTCAATGAOGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAACTCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTGCAACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 25 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

or an equivalent thereof. coding for said envelope protein fragment.

32. An expression vector according to claim 29, wherein said gene coding for an envelope protein fragment of an AIDS virus is a gene comprising the nucleotide sequence:

ATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAACAGATAGCTAGC
 35 AAATTAAGAGAACAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACGCACAGTTTTAATTGTGGAGGGGAATTTTCTACTGTAATTCACACAACCTGTTTAATAGT
 ACTTGGTTTAATAGTACTTGGAGTACTGAACGGTCAAATAACACTGAAOGAAGTGACACAATCACACTC
 CCATGCAGAAATAAACCAATTTATAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAAACAAGAGATGGTGGTAATAACAAC
 40 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTTGAACCAATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGAOGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 45 GCAAGAACTCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTGCAACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

50 or an equivalent thereof. coding for said envelope protein fragment.

33. An expression vector according to claim 29, wherein said gene coding for an envelope protein fragment of an AIDS virus is a gene comprising the nucleotide sequence:

55

ATGTATGCCCCCTCCCATC

AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 5 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTTGGGTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 10 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCCTTGAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

or an equivalent thereof. coding for said envelope protein fragment.

34. An expression vector according to claim 29, wherein said gene coding for an envelope protein fragment of an AIDS virus is a gene comprising the nucleotide sequence:

ATGAGGGACAATTGGAGAAGTGAATTATATAAA

TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTTGGGTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAAT
 25 TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCCTTGAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

35. An expression vector according to any one of claims 29 to 34 which is a plasmid capable of replication in gram-negative bacteria.

36. An expression vector according to claim 35 which is capable of replication in an E. coli strain.

37. The expression vector pEV1, -2, or -3/env 44-640.

38. The expression vector pEV1, -2, or -3/env 205-640.

39. A transformant carrying an expression vector as claimed in any one of claims 29-38.

40. A transformant according to claim 39 which is an E. coli strain.

41. A transformant according to claim 40 which is an E. coli MC 1061 strain.

42. Antibodies raised against a protein fragment obtained according to claims 1 to 5 and 26.

43. The antibodies of claim 42 which are monoclonal antibodies.

44. A vaccine eliciting immunity to AIDS comprising as an active ingredient a protein fragment obtained according to claims 1 to 5 and 26.

45. The use of a protein fragment as claimed in claim 1 for the preparation of a protective immunisation vaccine.

Patentansprüche

Patentansprüche für folgend Vertragsstaaten : BE, CH, DE, FR, GB, IT, LI, NL, SE,

- 5 1. Ein Hüllproteinfragment eines Erworbenen-Immunschwäche-Syndrom-(AIDS)-Virus, weitgehend frei von anderen Proteinen, mit der Aminosäuresequenz:

ValTrpLysGluAla
 10 ThrThrThrLeuPheCysAlaSerAspAlaLysAlaTyrAspThrGluValHisAsnValTrpAlaThr
 HisAlaCysValProThrAspProAsnProGlnGluValValLeuValAsnValThrGluAsnPheAsn
 METTrpLysAsnAspMETValGluGlnMETHisGluAspIleIleSerLeuTrpAspGlnSerLeuLys
 ProCysValLysLeuThrProLeuCysValSerLeuLysCysThrAspLeuLysAsnAspThrAsnThr
 15 AsnSerSerSerGlyArgMETIleMETGluLysGlyGluIleLysAsnCysSerPheAsnIleSerThr
 SerIleArgGlyLysValGlnLysGluTyrAlaPhePheTyrLysLeuAspIleIleProIleAspAsn
 AspThrThrSerTyrThrLeuThrSerCysAsnThrSerValIleThrGlnAlaCysProLysValSer
 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 20 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 25 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 30 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 35 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

40 oder

45

50

55

CysProLysValSer

PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 5 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 10 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 15 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 20 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

oder

METArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 30 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 35 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 40 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

oder

METTyrAlaProProIle

SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 50 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 55 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

TGTCCAAAGGTATCC

TTTGAGCCCAATTCCTATACATTAATGTCGCCCCGGCTGGTTTTGCGATTCTAAAAATGTAATAAAGACG
 5 TTTCAATGGAAACAGGACCATGTACAAAATGTCAGGCACAGTACAAATGTACACATGGAAATTAGGCCAGTAGTA
 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAGAAAGAGGTAGTAATTAGATCTGTCAATTTTCAGG
 GACAAATGCTAAAACCATTAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCCAACAC
 AATACAAGAAAAAAATCCGTATCCAGAGGGGACCCAGGGAGAGCATTGTTTACAAATAGGAAAAATAGGA
 AATATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAAATGCCACTTTAAACAGATAGCTAGC
 10 AAATTAAGAGAAACAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAAAGCACAGTTTTAAATTTGAGAGGGGAATTTTCTACTGTAAATTCACACAACTGTTTTAATAGT
 ACTTGGTTTTAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCCAGAATAAAACAATTTATAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAAATAACAAC
 15 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTCGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATGAAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTCTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTAATGTTCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 20 GCAAGAACTCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGAATTTGGGGTTGCTCTGGA
 AAACATAATTCACCACTGCTGTGCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAAACAGATTTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

oder ein Äquivalent davon kodierend für das besagte Hüllproteinfragment.

5. Ein Expressionsvektor gemäss Anspruch 2, worin das besagte, für ein Hüllproteinfragment eines AIDS-Virus kodierende Gen ein Gen ist, das die folgende Nukleinsäuresequenz enthält:

ATGAGACAAGCACATTGTAAACATTAGTAGAGCAAAATGGAAATGCCACTTTAAACAGATAGCTAGC
 AAATTAAGAGAACAAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAAAGCACAGTTTTAAATTTGAGAGGGGAATTTTCTACTGTAAATTCACACAACTGTTTTAATAGT
 ACTTGGTTTTAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACACTC
 35 CCATGCCAGAATAAAACAATTTATAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTCGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATGAAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTCTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 40 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTAATGTTCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAACTCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGAATTTGGGGTTGCTCTGGA
 AAACATAATTTGCACCACTGCTGTGCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAAACAGATTTGG
 45 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

oder ein Äquivalent davon kodierend für das besagte Hüllproteinfragment.

6. Ein Expressionsvektor gemäss Anspruch 2, worin das besagte, für ein Hüllproteinfragment eines AIDS-Virus kodierende Gen ein Gen ist, das die folgende Nukleinsäuresequenz enthält:

5
 10
 15
 20
 25
 30

ATGTATGCCCCCTCCCATC
 AGGGGACAAATTAGATGTTCTCAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGA CAATTGGAGAAGTGAAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACCGCTGACGGTACAGGCCAGACAATTATTGTTCTGGTATAGTGCAGCAGCAGAACCAAT
 TTGCTGAGGGCTATTGAGGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AACTAAATTTGACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAAATAATCTCTGGAACAGATTGG
 AATCACCGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

oder ein Äquivalent davon kodierend für das besagte Hüllproteinfragment.

7. Ein Expressionsvektor gemäss Anspruch 2, worin das besagte, für ein Hüllproteinfragment eines AIDS-Virus kodierende Gen ein Gen ist, das die folgende Nukleinsäuresequenz enthält:

20
 25
 30

ATGAGGGACAAATTGGAGAAGTGAAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACCGCTGACGGTACAGGCCAGACAATTATTGTTCTGGTATAGTGCAGCAGCAGAACCAAT
 TTGCTGAGGGCTATTGAGGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AACTAAATTTGACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAAATAATCTCTGGAACAGATTGG
 AATCACCGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

- 35
 40
 45
 50
 55
8. Ein Expressionsvektor gemäss einem der Ansprüche 2 bis 7, der ein Plasmid ist, das sich in gram-negativen und/oder gram-positiven Bakterien replizieren kann.
 9. Ein Expressionsvektor gemäss Anspruch 8, welcher fähig ist, in einem E. coli Stamm zu replizieren.
 10. Ein Expressionsvektor gemäss Anspruch 8, welcher fähig ist, in einem B. subtilis Stamm zu replizieren.
 11. Der Expressionsvektor pEV1, -2, oder -3/env 44-640.
 12. Der Expressionsvektor pEV1, -2, oder -3/env 205-640.
 13. Ein Transformant der einen Expressionsvektor gemäss einem der Ansprüche 2 bis 12 trägt.
 14. Ein Transformant gemäss Anspruch 13, der ein E. coli Stamm ist.
 15. Ein Transformant gemäss Anspruch 13, der ein E. coli MC 1061 Stamm ist.
 16. Ein Transformant gemäss Anspruch 13, der ein B. subtilis Stamm ist.
 17. Ein Transformant gemäss Anspruch 13, welcher eine Säugetierzelle ist.
 18. Ein Verfahren zur Herstellung eines wie in Anspruch 1 beanspruchten Hüllproteinfragments eines Erworbenen-Immunschwäche-Syndrom-Virus gekennzeichnet durch:

Transformieren einer Wirtszelle mit einem Expressionsvektor wie in einem der Ansprüche 2 bis 12 beansprucht; Kultivieren besagter Wirtszelle, so dass besagtes AIDS env Proteinfragment exprimiert wird;

und Extrahieren und Isolieren des besagten AIDS env Proteinfragments.

19. Ein Verfahren gemäss Anspruch 18, worin der Expressionsvektor pEV1, -2 oder -3/env 44-640 ist.

5 20. Ein Verfahren gemäss Anspruch 18, worin der Expressionsvektor pEV1, -2 oder -3/env 205-640 ist.

21. Ein Verfahren zum Testen von humanem Blut auf das Vorhandensein des viralen Verursachers von AIDS, gekennzeichnet durch Mischen einer Zusammensetzung enthaltend ein Hüllproteinfragment eines AIDS Virus gemäss Anspruch 1 mit einer Probe von humanem Blut und Bestimmen ob das besagte Hüllproteinfragment an in der Blutprobe vorhandene AIDS Antikörper bindet.

22. Ein Verfahren gemäss Anspruch 21, gekennzeichnet durch die Verwendung der Western Blot Analyse umfasst.

15 23. Ein Verfahren gemäss Anspruch 21, gekennzeichnet durch die Verwendung einer ELISA Technik, wobei ein Hüllproteinfragment eines AIDS Virus gemäss Anspruch 1 auf eine Festphase aufgebracht wird, mit der Probe in Kontakt gebracht wird und nach Waschen mit einem enzymmarkiertem nicht-humanem IgG zusammengebracht wird.

24. Ein Verfahren gemäss Anspruch 21, worin das Doppel-Antigen-Verfahren verwendet wird.

20 25. Ein Verfahren zur Bestimmung von AIDS-Viren, worin Antikörper gegen das Hüllproteinfragment eines AIDS-Virus gemäss Anspruch 1 verwendet werden.

26. Ein Verfahren gemäss Anspruch 25, worin das Antigen in der Probe und ein Proteinfragment gemäss Anspruch 1 welches markiert ist, um einen Antikörper gegen ein Proteinfragment gemäss Anspruch 1 konkurrieren.

25 27. Ein Verfahren gemäss Anspruch 25, worin ein Sandwichverfahren unter Verwendung von zwei Antikörpern gegen ein Proteinfragment gemäss Anspruch 1 durchgeführt wird.

30 28. Ein Verfahren gemäss Anspruch 27, worin ein Antikörper an der Festphase ist und der andere Antikörper markiert ist.

29. Ein Verfahren gemäss Anspruch 27, worin zwei verschiedene monoklonale Antikörper verwendet werden.

35 30. Ein Immunität gegen AIDS bewirkender Impfstoff, enthaltend als aktiven Bestandteil ein Proteinfragment gemäss Anspruch 1.

31. Antikörper erzeugt gegen ein Proteinfragment gemäss Anspruch 1.

40 32. Die Antikörper gemäss Anspruch 31, welche monoklonale Antikörper sind.

33. Die Verwendung eines Proteinfragments gemäss Anspruch 1 für die Herstellung eines schützenden immunisierenden Impfstoffs.

45 34. Die Verwendung eines Proteinfragments gemäss Anspruch 1 zum Testen von humanem Blut auf das Vorhandensein von AIDS-Viren.

Patentansprüche für folgenden Vertragsstaat : AT

50 1. Verfahren für die Herstellung eines Hüllproteinfragments eines Erworbenen-Immunschwäche-Syndrom-(AIDS)-Virus, welches im wesentlichen frei von anderen Proteinen ist, gekennzeichnet durch:

Transformieren einer Wirtszelle mit einem Expressionsvektor enthaltend ein Gen kodierend für ein Hüllproteinfragment eines AIDS-Virus mit der Aminosäuresequenz:

55

ValTrpLysGluAla
 ThrThrThrLeuPheCysAlaSerAspAlaLysAlaTyrAspThrGluValHisAsnValTrpAlaThr
 HisAlaCysValProThrAspProAsnProGlnGluValValLeuValAsnValThrGluAsnPheAsn
 5 METTrpLysAsnAspMETValGluGlnMETHisGluAspIleIleSerLeuTrpAspGlnSerLeuLys
 ProCysValLysLeuThrProLeuCysValSerLeuLysCysThrAspLeuLysAsnAspThrAsnThr
 AsnSerSerSerGlyArgMETIleMETGluLysGlyGluIleLysAsnCysSerPheAsnIleSerThr
 SerIleArgGlyLysValGlnLysGluTyrAlaPhePheTyrLysLeuAspIleIleProIleAspAsn
 10 AspThrThrSerTyrThrLeuThrSerCysAsnThrSerValIleThrGlnAlaCysProLysValSer
 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 SerThrGlnLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 15 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 20 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 25 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 30 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

oder

CysProLysValSer
 35 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 40 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 45 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 50 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 55 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

oder

METArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnS rSerGlyGlyAspProGlu
 5 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 10 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 15 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

oder

20 METTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 25 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 30 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

oder

35 METArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 40 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 45 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

50 abwärts von einer Promotorsequenz, die die Transkription, Translation und damit die Expression des
 Hüllproteinfragments in einer Wirtszelle ermöglicht; Kultivieren der Wirtszelle, so dass das Hüllproteinfragment
 eines AIDS-Virus expremiert wird; und Extrahieren und Isolieren des Hüllproteinfragments von einem AIDS-
 Virus.

2. Ein Verfahren gemäss Anspruch 1, worin die Wirtszelle ein Bakterium ist.
3. Ein Verfahren gemäss Anspruch 2, worin das Bakterium E. coli ist.
4. Ein Verfahren gemäss Anspruch 3, worin das Plasmid pEV1, -2 oder 3/env 44-640 ist.
5. Ein Verfahren gemäss Anspruch 3, worin das Plasmid pEV1, -2 oder 3/env 205-640 ist.

6. Ein Verfahren für die Herstellung eines Expressionsvektors enthaltend ein Gen kodierend für ein Hüllproteinfragment eines AIDS-Virus, gekennzeichnet durch das Konstruieren eines Expressionsvektors mit einer Inserierungsstelle, worin das in Anspruch 1 definierte Gen kodierend für ein Hüllproteinfragment eines AIDS-Virus inseriert werden kann, wobei die Inserierungsstelle aufwärts einer Promotorsequenz liegt, die die Transkription, Translation und damit Expression des Hüllproteinfragments in einer Wirtszelle ermöglicht.
7. Ein Verfahren gemäss Anspruch 6, dadurch gekennzeichnet, dass als Gen, welches für ein Hüllproteinfragment eines AIDS-Virus kodiert, ein Gen enthaltend die Nukleotidsequenz:

GTGTGGAAGGAAGCA
 ACCACCACTCTATTTTGTGCATCAGATGCTAAAGCATATGATACAGAGGTACATAATGTTTGGGCCACA
 CATGCCTGTGTACCCACAGACCCCAACCCCAAGAAGTAGTATTGGTAAATGTGACAGAAAAATTTTAAC
 ATGTGGAATAATGACATGGTAGAACAGATGCATGAGGATATAATCAGTTTATGGGATCAAAGCCTAAAG
 CCATGTGTAAAAATTAACCCCACTCTGTGTAGTTTAAAGTGCACTGATTTGAAGAATGATACTAATACC
 AATAGTAGTAGCGGGAGAATGATAATGGAGAAAGGAGAGATAAAAACTGCTCTTTCAATATCAGCACA
 AGCATAAGAGGTAAAGGTGCAGAAAGAAATATGCATTTTTTTTAAACTTGATATAATACCAATAGATAAT
 GATACTACCAGCTATACGTTGACAGTTGTAAACCTCAGTCATTACACAGGCCTGTCCRAAGGTATCC
 TTTGAGCCAAATCCCATACATTATTTGTGCCCCGGCTGGTTTTGCGATTCTAAAAATGTAATAATAAGAG
 TTCAATGGAACAGGACCATGTACAAATGTCAGCACAGTACAATGTACACATGGAATTAGGCCAGTAGTA
 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAAGAGGTAGTAATTAGATCTGTCAATTTTCAAG
 GACAATGCTAAAAACCATAAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCACAAC
 AATCAAGAAAAAAATCCGTATCCAGAGGGGACCAGGGAGAGCATTGTTACAAATAGGAAAAATAGGA
 AATATGAGACAAGCATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAACAGATAGCTAGC
 AAATTAAGAGAACAATTTGGAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGACCCAGAA
 ATTGTAACGCACAGTTTTTAATTGTGGAGGGGAATTTTTCTACTGTAATTCACACAACCTGTTTAATAGT
 ACTTGGTTTAAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCAGAATAAAACAATTTATAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCCCTCCCATC
 AGCGGACAAATTAGATGTTCTCRAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACRAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATRAA
 TATAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCRAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTTCTGGTATAGTGACGACAGCAGAACAT
 TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCTAG
 GCAAGAAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTTGG
 AATCACACCAGTGGATGGAGTGGGACAGAGAAATTAACAAATTACACAAGC

oder ein dafür kodierendes Aequivalent verwendet wird.

8. Ein Verfahren gemäss Anspruch 6, dadurch gekennzeichnet, dass als Gen, welches für ein Hüllproteinfragment eines AIDS-Virus kodiert, ein Gen enthaltend die Nukleotidsequenz:

TGTCCAAAGGTATCC

TTTGAGCCAATTCCCATACATTATTGTGCCCCGGCTGGTTTTGCGATTCTAAATGTAATAAAGAGG
 TTCAATOGAACAGGACCATGTACAAATGTCAGCACAGTACAATGTACACATGGAATTAGGCCAGTAGTA
 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAAGAGGTAGTAATTAGATCTGTCAATTTACAG
 GACAACTCTAAACCATAATAGTACAGCTGAACACATCTGTAGAAATTAATGTACAAGACCCACACAC
 AATACAAGAAAAAAATCCGTATCCAGAGGGGACCAGGGAGAGCATTGTTTACAATAGGAAAAATAGGA
 AATATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAACAGATAGCTAGC
 AAATTAAGAGAACAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACGCACAGTTTTAATTGTGGAGGGGAATTTTTCTACTGTAATTCACACAACCTGTTTAATAGT
 ACTTGGTTTAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCAGAATAAAACAATTTATAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTGGGTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAACATAATTCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTTGG
 AATCACACGACGTGGATGGAGTGGACAGAGAAATTAACAATTACACAAGC

oder ein dafür kodierendes Aequivalent verwendet wird.

9. Ein Verfahren gemäss Anspruch 6, dadurch gekennzeichnet, dass als Gen, welches für ein Hüllproteinfragment eines AIDS-Virus kodiert, ein Gen enthaltend die Nukleotidsequenz:

ATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAACAGATAGCTAGC
 AAATTAAGAGAACAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACGCACAGTTTTAATTGTGGAGGGGAATTTTTCTACTGTAATTCACACAACCTGTTTAATAGT
 ACTTGGTTTAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCAGAATAAAACAATTTATAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTGGGTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAACATAATTCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTTGG
 AATCACACGACGTGGATGGAGTGGACAGAGAAATTAACAATTACACAAGC

oder ein dafür kodierendes Aequivalent verwendet wird.

10. Ein Verfahren gemäss Anspruch 6, dadurch gekennzeichnet, dass als Gen, welches für ein Hüllproteinfragment eines AIDS-Virus kodiert, ein Gen enthaltend die Nukleotidsequenz:

ATGTATGCCCTCCCATC

AGCGGACAAATTAGATGTTCTCAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 5 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGCCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCCTGGTATAGTGCAGCAGCAGAACRAAT
 TTGCTGAGGGCTATTGAGGGCGCAACAGCATCTGTTGCAACTCAAGTCTGGGGCATCAGCAGCTCCAG
 10 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

15 oder ein dafür kodierendes Aequivalent verwendet wird.

11. Ein Verfahren gemäss Anspruch 6, dadurch gekennzeichnet, dass als Gen, welches für ein Hüllproteinfragment eines AIDS-Virus kodiert, ein Gen enthaltend die Nukleotidsequenz:

ATGAGGGACAATTGGAGAAGTGAATTATATAAA

TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGCCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCCTGGTATAGTGCAGCAGCAGAACRAAT
 25 TTGCTGAGGGCTATTGAGGGCGCAACAGCATCTGTTGCAACTCAAGTCTGGGGCATCAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGCT

30 oder ein dafür kodierendes Aequivalent verwendet wird.

12. Ein Verfahren gemäss einem der Ansprüche 6 bis 11, worin der Expressionsvektor ein Plasmid ist, das zur Replikation in gram-negativen Bakterien fähig ist.
- 35 13. Ein Verfahren gemäss Anspruch 12, worin das Plasmid zur Replikation in einen E.coli Stamm fähig ist.
14. Ein Verfahren für die Herstellung eines Transformanten, der einen Expressionsvektor enthaltend ein Gen kodierend für ein Hüllproteinfragment eines AIDS-Virus trägt, welches Verfahren Transformieren eines Mikroorganismus mit einem Expressionsvektor gemäss einem der Ansprüche 6 bis 13 und Kultivieren des transformierten Mikroorganismus umfasst.
- 40 15. Ein Verfahren gemäss Anspruch 14, worin der Mikroorganismus ein E.coli Stamm ist.
- 45 16. Ein Verfahren gemäss Anspruch 15, worin der Mikroorganismus eine E. coli MC 1061 Stamm ist.
17. Ein Verfahren zum Testen von humanem Blut auf das Vorhandensein des viralen Verursachers von AIDS, gekennzeichnet durch Mischen einer Zusammensetzung enthaltend ein Hüllproteinfragment eines AIDS-Virus erhalten gemäss Anspruch 1 mit einer Probe von humanem Blut und Bestimmen, ob das Hüllproteinfragment an in der Blutprobe vorhandene AIDS Antikörper bindet.
- 50 18. Ein Verfahren gemäss Anspruch 17, gekennzeichnet durch die Verwendung der Western Blot Analyse.
19. Ein Verfahren gemäss Anspruch 17, gekennzeichnet durch die Verwendung einer ELISA-Technik, wobei ein Hüllproteinfragment eines AIDS-Virus erhalten gemäss Anspruch 1 auf eine Festphase aufgebracht, mit der Probe in Kontakt gebracht und nach Waschen mit einem enzymmarkierten nicht-humanem IgG zusammengebracht wird.
- 55 20. Verfahren gemäss Anspruch 17, worin die Doppel-Antigen-Methode verwendet wird.

21. Ein Verfahren zur Bestimmung von AIDS-Viren, worin Antikörper gegen das gemäss Anspruch 1 erhaltene Hüllproteinfragment eines AIDS-Virus verwendet werden.
- 5 22. Ein Verfahren gemäss Anspruch 21, worin das Antigen in der Probe und ein Proteinfragment erhalten gemäss Anspruch 1, welches markiert ist, um einen Antikörper gegen ein Proteinfragment erhalten gemäss Anspruch 1 konkurrieren.
23. Ein Verfahren gemäss Anspruch 21, worin ein Sandwichverfahren unter Verwendung von zwei Antikörpern gegen ein gemäss Anspruch 1 erhaltenes Proteinfragment durchgeführt wird.
- 10 24. Ein Verfahren gemäss Anspruch 23, worin ein Antikörper an der Festphase ist und der andere Antikörper markiert ist.
25. Ein Verfahren gemäss Anspruch 23, worin zwei verschiedene monoklonale Antikörper verwendet werden.
- 15 26. Ein Hüllproteinfragment von einem AIDS-Virus, hergestellt durch ein Verfahren gemäss einem der Ansprüche 1 bis 5.
27. Ein Expressionsvektor, enthaltend ein Gen kodierend für ein Hüllproteinfragment eines AIDS-Virus, hergestellt durch ein Verfahren gemäss einem der Ansprüche 6 bis 13.
- 20 28. Ein Transformant tragend einen Expressionsvektor enthaltend ein Gen kodierend für ein Hüllproteinfragment eines AIDS-Virus, hergestellt durch ein Verfahren gemäss einem der Ansprüche 14 bis 16.
- 25 29. Ein Expressionsvektor enthaltend ein Gen kodierend für ein Hüllproteinfragment von einem AIDS-Virus gemäss Anspruch 1, abwärts von einer Promotorsequenz, die die Transkription, Translation und damit die Expression des besagten Hüllproteinfragments in einer Wirtszelle ermöglicht.
- 30 30. Ein Expressionsvektor gemäss Anspruch 29, worin das für ein Hüllproteinfragment eines AIDS-Virus kodierende Gen ein Gen ist, das die Nukleotidsequenz:

GTGTGGAAGGAAGCA

ACCACCACTCTATTTTGTGCATCAGATGCTAAAGCATATGATACAGAGGTACATAATGTTTGGGCCCA
 CATGCCCTGTGTACCCACAGACCCCAACCCCAAGAAGTAGTATTGGTAAATGTGACAGAAAATTTTAC
 5 ATGTGGA AAAATGACATGGTAGAACAGATGCATGAGGATATAATCAGTTTATGGGATCAAAGCCTAAG
 CCATGTGTAAAATTAACCCCACTCTGTGTTAGTTTAAAGTGCACTGATTTGAAGAAATGATACTAATACC
 AATAGTAGTAGCGGGAGAATGATAATGGAGAAAGGAGAGATAAAAAAAGTCTCTTTCAATATCAGCACA
 AGCATRAGAGGTAAAGGTGCAGAAAGAATATGCATTTTTTTTATAAACTTGATATAATACCAATAGATAAT
 10 GATACTACCAGCTATACGTTGACAAGTTGTAACACCTCAGTCATTACACAGGCCTGTCCAAAGGTATCC
 TTTGAGCCAAATCCCATACATTATTGTGCCCCGGCTGGTTTTGCGATTCTRAAATGTAATAAAGACG
 TTCAATGGAACAGGACCATGTACAAATGTCAGCACAGTACAAATGTACACATGGAATTAGGCCAGTAGTA
 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAAGAGGTAGTAATTAGATCTGTCAATTTCAAG
 GACAAATGCTAAAACCATAAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACRAGACCCACAAC
 15 AATACAGAAAAAAATCCGTATCCAGAGGGGACAGGGAGAGCAATTTGTTACAATAGGAAAAATAGGA
 AATATGAGACACAGCATTGTAACTTAGTAGAGCAAAATGGAAATGCCACTTTAAACAGATAGCTAGC
 AAATTAAGAGAACAAATTTGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAAACGACAGTTTAAATTTGTGGAGGGGAATTTTTCTACTGTAATTCRACACAACCTGTTAATAGT
 ACTTGGTTTAAATAGTACTTGGAGTACTGAAGGGTCRAATAACACTGAAGGAAGTGACCAATCACACTC
 20 CCRTGCAGAAATAAAACAATTTATAAACHTGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTATCAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAGTAGTAAAAATGAACCATTAGGAGTAGCAACCCACRAGGCAAAGAGAAGAGTGGTGACAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 25 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAACTAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 30 AATCACACGAGCTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

oder ein Äquivalent davon kodierend für das besagte Hüllproteinfragment enthält.

- 35 31. Ein Expressionsvektor gemäss Anspruch 29, worin das besagte, für ein Hüllproteinfragment eines AIDS-Virus kodierende Gen ein Gen ist, das die Nukleotidsequenz:

40

45

50

55

TGTCCAAAGGTATCC

TTTGAGCCAAATCCCATACATTATTGTGCCCGGCTGGTTTTGCGATTCTAAAAATGTAATAAAGACG
 TTCAATGGAACAGGACCATGTACAAATGTCAGCACAGTACAAATGTACACATGGAAATAGGCCAGTAGTA
 5 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAAGAGGTAGTAATTAGATCTGTCAATTTCCAG
 GACAATGCTAAAACCATTAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCCAAC
 AATACAAGAAAAAAATCCGTATCCAGAGGGGACCAGGAGAGCATTGTTACAAATAGGAAAAATAGGA
 AATATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAAACAGATAGCTAGC
 10 AAATTAAGAGAACAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCGAA
 ATTGTAACGCACAGTTTTAATTGTGGAGGGGAATTTTCTACTGTAATTCACACAACTGTTTAAATAGT
 ACTTGGTTTAAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCAGAATAAAACAATTTATAAACAATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCCCTCCATC
 AGCGGACAAATTAGATGTTTCAATCAATATTACAGGGCTGCTATTAAACAAGAGATGGTGGTAATAACAC
 15 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAA
 TATAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTGGGTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGGCTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAT
 TTGCTGAGGGCTATTGAGGGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 20 GCAAGAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAACATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

oder ein Äquivalent davon kodierend für das besagte Hüllproteinfragment enthält.

32. Ein Expressionsvektor gemäss Anspruch 29, worin das für ein Hüllproteinfragment eines AIDS-Virus kodierende Gen ein Gen ist, das die Nukleotidsequenz:

ATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAATGCCACTTTAAAACAGATAGCTAGC
 AAATTAAGAGAACAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCGAA
 ATTGTAACGCACAGTTTTAATTGTGGAGGGGAATTTTCTACTGTAATTCACACAACTGTTTAAATAGT
 ACTTGGTTTAAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACACTC
 35 CCATGCAGAATAAAACAATTTATAAACAATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCCCTCCATC
 AGCGGACAAATTAGATGTTTCAATCAATATTACAGGGCTGCTATTAAACAAGAGATGGTGGTAATAACAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAA
 TATAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTGGGTCTTGGGAGCAGCAGGAAGCACTATGGGC
 40 GCAGGCTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAT
 TTGCTGAGGGCTATTGAGGGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAACATAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTTGG
 45 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

oder ein Äquivalent davon kodierend für das besagte Hüllproteinfragment enthält.

33. Ein Expressionsvektor gemäss Anspruch 29, worin das besagte, für ein Hüllproteinfragment eines AIDS-Virus kodierende Gen ein Gen ist, das die Nukleotidsequenz:

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55

ATGTATGCCCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCGGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAT
 TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAACATAATTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGAGCTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

oder ein Äquivalent davon kodierend für das besagte Hüllproteinfragment enthält.

34. Ein Expressionsvektor gemäss Anspruch 29, worin das für ein Hüllproteinfragment eines AIDS-Virus kodierende Gen ein Gen ist, das die Nukleotidsequenz:

20
 25
 30
 35
 40
 45
 50
 55

ATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAT
 TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAACATAATTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGAGCTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

35. Ein Expressionsvektor gemäss einem der Ansprüche 29 bis 34, der ein Plasmid ist, das sich in gram-negativen Bakterien replizieren kann.

36. Ein Expressionsvektor gemäss Anspruch 35, welcher fähig ist, in einen E. coli Stamm zu replizieren.

37. Der Expressionsvektor pEV1, -2, oder -3/env 44-640.

38. Der Expressionsvektor pEV1, -2, oder -3/env 205-640.

39. Ein Transformant der einen Expressionsvektor gemäss einem der Ansprüche 29 bis 38 trägt.

40. Ein Transformant gemäss Anspruch 39, der ein E. coli Stamm ist.

41. Ein Transformant gemäss Anspruch 40, der ein E. coli MC 1061 Stamm ist.

42. Antikörper erzeugt gegen ein wie gemäss Ansprüchen 1 bis 5 und 26 erhaltenes Proteinfragment.

43. Die Antikörper von Anspruch 42, welche monoklonale Antikörper sind.

44. Ein Impfstoff der Immunität gegen AIDS bewirkt, enthaltend als aktiven Bestandteil ein Proteinfragment erhalten gemäss Ansprüchen 1 bis 5 und 26.

45. Die Verwendung eines wie in Anspruch 1 beanspruchten Proteinfragments zur Herstellung eines schützenden, immunisierenden Impfstoffes.

Revendications

Revendications pour les Etats contractants suivants : BE, CH, DE, FR, GB, IT, LI, NL, SE

- 5 1. Fragment d'une protéine d'enveloppe d'un virus du syndrome de l'immunodéficience acquise (SIDA), pratiquement exempté d'autres protéines, ayant la séquence d'acides aminés suivante :

ValTrpLysGluAla
 10 ThrThrThrLeuPheCysAlaSerAspAlaLysAlaTyrAspThrGluValHisAsnValTrpAlaThr
 HisAlaCysValProThrAspProAsnProGlnGluValValLeuValAsnValThrGluAsnPheAsn
 METTrpLysAsnAspMETValGluGlnMETHisGluAspIleIleSerLeuTrpAspGlnSerLeuLys
 ProCysValLysLeuThrProLeuCysValSerLeuLysCysThrAspLeuLysAsnAspThrAsnThr
 15 AsnSerSerSerGlyArgMETIleMETGluLysGlyGluIleLysAsnCysSerPheAsnIleSerThr
 SerIleArgGlyLysValGlnLysGluTyrAlaPhePheTyrLysLeuAspIleIleProIleAspAsn
 AspThrThrSerTyrThrLeuThrSerCysAsnThrSerValIleThrGlnAlaCysProLysValSer
 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 20 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 25 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 30 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 35 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

ou

40

45

50

55

CysPr LysValSer

PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 5 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 10 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 15 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 20 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

OU

25

METArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 30 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 35 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 40 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

OU

45

METTyrAlaProProIle

SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 50 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

55

OU

METArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer.

2. Vecteur d'expression comprenant un gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA telle que définie dans la revendication 1 en aval d'une séquence de promoteur permettant la transcription, la traduction et, par conséquent, l'expression de ce fragment de cette protéine d'enveloppe dans une culture hôte.
3. Vecteur d'expression selon la revendication 2, dans lequel ce gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA est un gène comprenant la séquence de nucléotides suivante :

GTGTGGAAGGAAGCA
 ACCACCACTCTATTTTGTGCATCAGATGCTAAAGCATATGATACAGAGGTACATAATGTTTGGGCCACA
 CATGCCCTGTGTACCCACAGACCCCAACCCACAAGAGTAGTATTGGTAAATGTGACAGAAAATTTTAAC
 ATGTGGAAAAATGACATCGTAGAACAGATGCATGAGGATATAATCAGTTTATCGGATCAAAGCCTAAAG
 CCATGTGTAAAAATTAACCCCACTCTGTGTAGTTTAAAGTGCCTGATTTGAAGAATGATACTAATACC
 AATAGTAGTAGCCGGAGAATGATAATGAGAGAAAGGAGAGATAAAAACTGCTCTTCAATATCAGCCACA
 AGCATAAGAGGTAAGGTGCAGAAAGAAATATGCATTTTTTATAAACTTGATATAATACCAATAGATAAT
 GATACTACCAGCTATACGTTGACAGTTGTAACACCTCAGTCATTACACAGGCCTGTCCAAAGGTATCC
 TTTGAGCCAAATCCCATACATTATGTGCCCCGGCTGGTTTTTGGGATTCTAAAATGTAATAAAGAGC
 TTCAATGGAAACAGGACCATGTACAAATGTCAGCACAGTACAATGTACACATCGAATTAGCCAGTAGTA
 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAAGAGGTAGTAATTAGATCTGTCAATTTACCG
 GACAATGCTAAAACCAATAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCAAACAAC
 AATACAAGAAAAAAATCCGTATCCAGAGGGGACCAGGGAGAGCATTGTTACAATAGGAAAAATAGGA
 AATATGAGACAAGCACAATTGTAACATTAGTAGACCAAAATGGAAATGCCACTTTAAACAGATAGCTAGC
 AAATTAAGAGAACAATTTGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACCCACAGCTTTTAATTTGCGAGGGGAATTTTTCTACTGTAATTCACACAACCTGTTAATAGT
 ACTTGGTTTAATAGTACTTGGAGTACTGAAGGGTCAAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCGAATAAAACAATTTATAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTCGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAAATTATATAAA
 TATAAAGTAGTAAAAATGAACCATTAGGAGTAGCACCCACCAAGGCAAGAGAAGAGTGGTGACAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGCTCGTATAGTGACGACGACAGAACAAT
 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAACTCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AAACATAATTTGCACCACTGCTGTGCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

ou un équivalent de celle-ci codant pour ledit fragment de la protéine d'enveloppe.

4. Vecteur d'expression selon la revendication 2, dans lequel ce gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA est un gène comprenant la séquence de nucléotides suivante :

TGTCCAAAGGTATCC

TTTGAGCCAATTCCCATACATTATTGTGCCCGGCTGGTTTTGCGATTCTAAAAATGTAATAAAGACG
 TTCAATGGAAACAGGACCATGTACAAAATGTCAGCACAGTACAATGTACACATCGAATTAGGCCAGTAGTA
 5 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAGAAGAGGTAGTAATTAGATCTGTCAATTTCCAG
 GACAAATGCTAAAAACCATAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCAACAAC
 AATACAAGAAAAAAATCCGTATCCAGAGGGGACCAGGAGAGCATTGTTACAATAGGAAAAATAGGA
 AATATGAGACAAAGCATTGTAAACATTAGTAGAGCAAAATGGAAATGCCACTTTAAACAGATAGCTAGC
 AAATTAAGAGAACAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGCGGACCCAGAA
 10 ATTGTAACGCACAGTTTTAATTGTGGAGGGGAATTTTTCTACTGTAATTCACACAACCTGTTTAATAGT
 ACTTGGTTTTAATAGTACTTGGAGTACTGAAGGGTCAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCAGAATAAAACAATTTATAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAAACAAGAGATCGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGACAAATGGAGAAGTGAATTATATAAA
 15 TATAAAGTAGTAAAAATTGAACCATAGGAGTAGCACCCACCAAGGCAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTCGGAATAGGAGCTTTGTTCTTGGGTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACCGTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAT
 TTGCTGAGGGCTATTGAGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTTGGGGTTGCTCTGGA
 20 AAATAATTTGCACCACTGCTGTGCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGACGTGGATGGAGTGGACAGAGAAATTAACAATTACACAAGC

ou un équivalent de celle-ci codant pour ledit fragment de la protéine d'enveloppe.

- 25 5. Vecteur d'expression selon la revendication 2, dans lequel ce gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA est un gène comprenant la séquence de nucléotides suivantes :

ATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAAATGCCACTTTAAAAACAGATAGCTAGC
 30 AAATTAAGAGAACAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACGCACAGTTTTAATTGTGGAGGGGAATTTTTCTACTGTAATTCACACAACCTGTTTAATAGT
 ACTTGGTTTTAATAGTACTTGGAGTACTGAAGGGTCAATAACACTGAAGGAAGTGACACAATCACACTC
 CCATGCAGAATAAAACAATTTATAAACATGTGCCAGGAAGTAGGAAAAGCAATGTATGCCCCCTCCCATC
 35 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAAACAAGAGATCGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGACAAATGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTCGGAATAGGAGCTTTGTTCTTGGGTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACCGTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAT
 40 TTGCTGAGGGCTATTGAGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTTGGGGTTGCTCTGGA
 AAATAATTTGCACCACTGCTGTGCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGG
 AATCACACGACGTGGATGGAGTGGACAGAGAAATTAACAATTACACAAGC

45 ou un équivalent de celle-ci codant parmi ledit fragment de la protéine d'enveloppe.

6. Vecteur d'expression selon la revendication 2, dans lequel le gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA est un gène comprenant la séquence de nucléotides suivante :

50

55

ATGTATGCCCTCCCATC

AGCGGACAAATTAGATGTTTCATCAAATATTACAGGGCTGCTATTAAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTCAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACCGTGACCGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAT
 TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCAEAGTCTGGGGCATCAAGCAGCTCCAG
 CCAAGAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTTGGGGTTGCTCTGGA
 AAACTAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAACTCTGGAACAGATTTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

ou un équivalent de celle-ci codant pour ledit fragment de la protéine d'enveloppe.

7. Vecteur d'expression selon la revendication 2, dans lequel ce gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA est un gène comprenant la séquence de nucléotides suivante :

ATGAGGGACAATTGGAGAAGTGAATTATATAAA

TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTCAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCCCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACCGTGACCGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAAT
 TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTTGGGGTTGCTCTGGA
 AAACTAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTTGGAGTAATAAACTCTGGAACAGATTTGG
 AATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

8. Vecteur d'expression selon l'une quelconque des revendications 2 à 7, qui est un plasmide capable de se répliquer dans des bactéries gram-négatives et/ou gram-positives.

9. Vecteur d'expression selon la revendication 8, qui est capable de se répliquer dans une souche d'E.coli.

10. Vecteur d'expression selon la revendication 8, qui est capable de se répliquer dans une souche de B.subtilis.

11. Vecteur d'expression pEV1, -2 ou -3/env. 44-640

12. Vecteur d'expression pEV1, -2 ou 3/env. 205-640.

13. Transformant portant un vecteur d'expression selon l'une quelconque des revendications 2 à 12.

14. Transformant selon la revendication 13, qui est une souche d'E.coli.

15. Transformant selon la revendication 14, qui est une souche d'E.coli MC 1061.

16. Transformant selon la revendication 13, qui est une souche de B.subtilis.

17. Transformant selon la revendication 13, qui est une cellule de mammifère.

18. Procédé de préparation d'un fragment d'une protéine d'enveloppe d'un virus du syndrome d'immunofciencia acquise selon la revendication 1, consistant à :

transformer une cellule hôte avec un vecteur d'expression selon l'une quelconque des revendications 2 à 12 ;
 cultiver cette cellule hôte de façon que ce fragment de cette protéine d'enveloppe du SIDA soit exprimée ; et

extraire et isoler ce fragment de cette protéine d'enveloppe du SIDA.

19. Procédé selon la revendication 19, dans lequel le vecteur d'expression est pEV1, -1, -2 ou -3/env.44-640
- 5 20. Procédé selon la revendication 19, dans lequel le vecteur d'expression est pEV1, -2 ou -3/env. 205-640.
21. Procédé de détection dans le sang humain de la présence d'anticorps pour l'agent étiologique viral du SIDA, qui consiste à mélanger une composition contenant un fragment d'une protéine d'enveloppe d'un virus du SIDA, selon la revendication 1, avec un échantillon de sang humain et de déterminer si ce fragment de cette protéine d'enve-
10 loppe du SIDA se lie aux anticorps du SIDA présents dans l'échantillon de sang.
22. Procédé selon la revendication 21 qui consiste à utiliser l'analyse par "Western Blotting".
23. Procédé selon la revendication 21 qui comprend l'utilisation d'une technique ELISA, dans laquelle un fragment
15 d'une protéine d'enveloppe d'un virus du SIDA, selon la revendication 1, est appliquée sur une phase solide et mise en contact avec l'échantillon et, après lavage, mise en contact avec une IgG non humaine marquée par une enzyme.
24. Procédé selon la revendication 21, dans lequel on utilise la Méthode du Double Antigène.
- 20 25. Procédé pour la détermination du virus du SIDA, dans lequel on utilise des anticorps contre un fragment d'une protéine d'enveloppe d'un virus du SIDA, selon la revendication 1.
26. Procédé selon la revendication 25, dans lequel l'antigène présent dans l'échantillon et un fragment d'une protéine
25 selon la revendication 1, sous forme marquée entrent en compétition avec un anticorps contre un fragment d'une protéine selon la revendication 1.
27. Procédé selon la revendication 25, dans lequel on applique une méthode sandwich en utilisant deux anticorps contre un fragment d'une protéine selon la revendication 1.
- 30 28. Procédé selon la revendication 27, dans lequel un anticorps est sur une phase solide et l'autre anticorps est marqué.
29. Procédé selon la revendication 27, dans lequel on utilise deux anticorps monoclonaux différents.
- 35 30. Vaccin déclenchant l'immunité au SIDA comprenant comme ingrédient actif un fragment d'une protéine selon la revendication 1.
31. Anticorps formés contre un fragment d'une protéine selon la revendication 1.
- 40 32. Anticorps selon la revendication 1, qui sont des anticorps monoclonaux.
33. Utilisation d'un fragment d'une protéine selon la revendication 1, pour la préparation d'un vaccin d'immunisation protectrice.
- 45 34. Utilisation d'un fragment d'une protéine selon la revendication 1 pour détecter dans le sang humain la présence du virus du SIDA.

Revendications pour l'Etat contractant suivant : AT

- 50 1. Procédé pour préparer un fragment d'une protéine d'enveloppe d'un virus du syndrome de l'immunodéficience acquise (SIDA), essentiellement exempt de autres protéines, qui consiste :

à transformer une cellule hôte avec un vecteur d'expression comprenant un gène codant pour un fragment
55 d'une protéine d'enveloppe d'un virus du SIDA ayant la séquence d'acides aminés suivante :

ValTrpLysGluAla

5 ThrThrThrLeuPheCysAlaSerAspAlaLysAlaTyrAspThrGluValHisAsnValTrpAlaThr
 HisAlaCysValProThrAspProAsnProGlnGluValValLeuValAsnValThrGluAsnPheAsn
 METTrpLysAsnAspMETValGluGlnMETHisGluAspIleIleSerLeuTrpAspGlnSerLeuLys
 ProCysValLysLeuThrProLeuCysValSerLeuLysCysThrAspLeuLysAsnAspThrAsnThr
 AsnSerSerSerOlyArgMETIleMETGluLysGlyGluIleLysAsnCysSerPheAsnIleSerThr
 SerIleArgOlyLysValGlnLysGluTyrAlaPhePheTyrLysLeuAspIleIleProIleAspAsn
 10 AspThrThrSerTyrThrLeuThrSerCysAsnThrSerValIleThrGlnAlaCysProLysValSer
 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 SerThrGlnLeuLeuLeuAsnOlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 15 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyOlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 20 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 25 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerOlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 30 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

OU

CysProLysValSer

35 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 40 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyOlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 45 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyOlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 50 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuOlyPheLeuGlyAlaAlaGlySerThrMETOly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 55 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

OU

5 METArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerOlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 10 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETTyrAlaProProIle
 SerOlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyOlyAsnAsnAsn
 AsnOlySerGluIlePheArgProGlyGlyOlyAspMETArgAspAsnTrpArgSerOluLeuTyrLys
 15 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETOly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerOlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpOlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuOlyIleTrpOlyCysSerGly
 20 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnIleThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

ou

20 METTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspOlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 25 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValOlyIleOlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerOly
 30 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnIleThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

ou

35 METArgAspAsnTrpArgSerOluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 40 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 45 AsnIleThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSer

50 en aval d'un promoteur permettant la transcription, la traduction et l'expression du fragment de cette protéine
 d'enveloppe dans la cellule hôte ; à cultiver cette cellule hôte de façon à exprimer le fragment de la protéine
 d'enveloppe d'un virus du SIDA ; et à extraire et à isoler le fragment de la protéine d'enveloppe d'un virus du
 SIDA.

2. Procédé selon la revendication 1, dans lequel la cellule hôte est une bactérie.
3. Procédé selon la revendication 2, dans lequel la bactérie est E. coli.
4. Procédé selon la revendication 3, dans lequel le plasmide est pEV1, -2 ou -3/env 44-640.
5. Procédé selon la revendication 3, dans lequel le plasmide est pEV1, -2 ou -3/env 205-640.

6. Procédé pour préparer un vecteur d'expression comprenant un gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA, procédé qui consiste à construire un vecteur d'expression portant un site d'insertion, dans lequel on peut insérer un gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA selon la revendication 1, le site d'insertion se trouvant en aval d'un promoteur permettant la transcription, la traduction et donc l'expression du fragment de la protéine d'enveloppe dans une cellule hôte.

7. Procédé selon la revendication 6, caractérisé en ce qu'on utilise en tant que gène codant pour un fragment d'une protéine d'enveloppe du virus du SIDA un gène comprenant la séquence nucléotidique suivante :

GTGTGGAAGGAAGCA

ACCACCCTCTATTTTGTGCATCAGATGCTAAAGCATATGATACAGAGGTACATAATGTTTGGGCCACA
 CATGCTGTGTACCCACAGACCCCAACCCACAAGAAGTACTATTGGTAAATGTGACAGAAATTTTAAC
 ATGTGGAAAAATGACATGGTAGACAGATGCATCAGGATATAATCAGTTTATGGGATCAAAOCCATAAG
 CCAITGTGTAAAAATTAACCCACTCTGTGTAGTTTAAAGTGCATGATTTGAAAGATGATACTAATACC
 AATACTAGTAGCCGGAGAAATGATAATGGAGAAAGAGAGATAAAAACTGCTCTTTCAATATCAGGCACA
 AGCATAGAGCGTAAGGTGCAGAAAGAAATATGCATTTTTTATAAACTTGTATATAATACCAATAGATAAT
 GATACTACCCAGCTATACGTTGACAAAGTTGTAACACCTCACTCATACACAGCCCTGTCCAAAGGTATCC
 TTTGAGCCAAATTCCTATACATTATTGTGCCCCCGCTGCTTTTGGGATTCTAAAAATGTAATAAAGACC
 TTCATGCAAGAGGACCATGTACAAATGTGAGCACAGTACAAATGTACACATGGAATTACGCCAGTAGTA
 TEAACTCAACTGCTTOTTAAAAAGCCAGTCTAGCAGAAAGAGGTAGTAATTAGATCTGTCAATTTACAG
 GACAAATGCT/AAAAACATAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCAAACAAC
 AATACAAGAAAAAAATCCOTATCCAGAGGGGACCCAGCAGCCATTTOTTACAATAGGAAAAATAGGA
 AATATGAGCAAGCACATTOTAACATTAGTAGAGCAAAATGCAATGCCACTTTAAAACAGATAOCTAGC
 AAAITTAGAGCAACPAATTTGGAATAAT/AAAAATTAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAACCCACAGTTTAAATTGTGACGGGGAATTTTCTACTGTAATTCACACAACCTOTTTAATAGT
 ACTTGGTTTAAATAGTACTTGGAGTACTGAAAGGTCAAAATACACTGAAGGAAGTGACACAATCACACTC
 CCATGCGAATAAAACAAATTTATAAACATGTGSCAGGAAGTAGGAAAAGCAATCTATGCCCTTCCCATC
 AGTGGACAAATTAGATGTTTATCAAAATATTACAGGCTGCTATTAAAGAGATGGTGGTAATAACAAC
 PATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGGACAAATGGAGAAGTGAATTATATAAA
 TATAAGTAGTAAAAATTAACCATTAGGAGTAGCACCCACCAAGGCAAGAGAGAGTGGTGCAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTTCTCCTTGGGTTCTTGGGAGCAGCAGGAGGCACTATGGGC
 GCACCCOTCAATGACCGTGACCGTACAGGCCAGACAATTATGTCTGGTATAGTGCAGCAGCAGAACAAT
 TTGCTGAGCGCTATTGAGCGGCAACAGCATCTGTTGCAACTCACAGTCTGGGSCATCAAGCAGCTCCAG
 GCAAGAACTCTGGCTGTGGAAAGATACCTAAAGGATGACAGCTCTGGCAATTTGGGTTGCTCTGGA
 AACTAATTTGCACCAGTGTGTGCTTGGAAATGCTAGTTGGAGTAAT/ATCTCTGACAGATTTCG
 AATCACACGACUTGGAATGGAGTGGGACAGAGAAATTAACAAATTACACAAGC

ou on utilise un équivalent codant en conséquence.

8. Procédé selon la revendication 6, caractérisé en ce qu'on utilise en tant que gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA un gène comprenant la séquence nucléotidique suivante :

TGTCCAAAGGTATCC
 TTTGAGCCAATTCCCATACATTATTGTGCCCCGGCTGCTTTTGGATTCTAAATGTAAATAAAGACO
 TTCAATGGAAACAGGACCATGTACAAAATGTCAGCACAGTACAAATGTACACATGGAAATTAGGCCAGTAGTA
 5 TCAACTCRACTGCTGTAAATGGCAGTCTAGCAGAAAGAGAGGTAGTAATTAGATCTGTCAATTTACCG
 GACAATGCTAAACCATAAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAGACCCAAACAC
 AATACAAGAAAAAAATCCGTATCCAGAGGGGACCAAGGAGAGCATTTTGTACAATAGAAAAATAGGA
 AATATGAGACAAGCACATTOTAACTTAGTAGAGCAAAATGGAAATGCCACTTTAAACACAGATAGCTAOC
 10 AAATTAAGAGAACAATTTGGAAATAATAAAACATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAAACGCACAGTTTTAATTGTGGAGGGGAATTTTCTACTGTAAATTCACACAACCTGTTTAAATAGT
 ACTTGGCTTTAATAGTACTTGGAGTACTGAAAGGGTCMAATAACACTGAAGGAAGTGACACAATCACACTC
 CCAATGCAGAAATAAAACAATTTATAAACATGTGGCAGGAAGTAAGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAACAAGAGATGGTGTAAATAACAAC
 15 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGACAATTCGAGAAGTGAATTAATAAA
 TATAAAGTAGTAAAAATTGAACCAATTAGGAGTAGCACCCACCAAGGCAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTCCCTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 GCAAGCTCAATGACCGTACCGGTACAGGCCAGACAATTAATGCTCTGTTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 20 GCAAGAACTCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGAAATTTGGGGTTGCTCTGGA
 AAATTAATTTGCACCACTGCTGTGCTTGGAAATGCTAGTTGGAGTAATAATCTCTGGAACAGATTGG
 AATCACACCAAGCTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAAC

25 ou on utilise un équivalent codant en conséquence.

9. Procédé selon la revendication 6, caractérisé en ce qu'on utilise comme gène codant pour un fragment d'une protéine d'enveloppe du virus du SIDA un gène comprenant la séquence nucléotidique suivante :

30 ATGAGACAAGCACATTGTAACTTAGTAGAGCAAAATGGAAATGCCACTTTAAACAGATAGCTAGC
 AATTAAGAGAACAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
 ATTGTAAACGCACAGTTTTAATTGTGGAGGGGAATTTTCTACTGTAAATTCACACAACCTGTTTAAATAGT
 ACTTGGCTTTAATAGTACTTGGAGTACTGAAAGGGTCMAATAACACTGAAGGAAGTGACACAATCACACTC
 35 CCAATGCAGAAATAAAACAATTTATAAACATGTGGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
 AGCGGACAAATTAGATGTTTCATCAAAATATTACAGGGCTGCTATTAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGACAATTCGAGAAGTGAATTAATAAA
 TATAAAGTAGTAAAAATTGAACCAATTAGGAGTAGCACCCACCAAGGCAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTCCCTTCTTGGGAGCAGCAGGAAGCACTATGGGC
 40 GCAAGCTCAATGACCGTACCGGTACAGGCCAGACAATTAATGCTCTGTTATAGTGCAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAACTCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGAAATTTGGGGTTGCTCTGGA
 AAATTAATTTGCACCACTGCTGTGCTTGGAAATGCTAGTTGGAGTAATAATCTCTGGAACAGATTGG
 45 AATCACACCAAGCTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAAC

ou on utilise un équivalent codant en conséquence.

10. Procédé selon la revendication 6, caractérisé en ce qu'on utilise comme gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA un gène comprenant la séquence nucléotidique suivante :

55

ATGTATGCCCTCCCATC

ACCGGACAAATTAGATGTTTCATCAAAATATTACAGCGCTCTATTAAACAAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGAGATATGAGGACAAATTCAGAAAGTGAATTATATAAA
 TATAAAGTATGTAATAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAGAGAGAGTGGTGCAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTCTGGGTCTTGGGAGCAGCAGGAAGCACTATGGC
 GCAGCGTCAATACGCTGACGGTACAGGCCAGACAATTATTGCTCTGGTATAGTCCAGCAGCAGAACAAT
 TTGCTGACGGCTATTGAGGCGAACAGCATCTGTTCCAACTCAGCTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAACTCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCTCGGGATTTGGGGTTGCTCTCGA
 AACTAATTTCCACCCTCTCTGTCCTTGGAACTGCTAGTTGGAGTAATAAATCTCTGGAACAGATTG
 AATCACACGACGTGGATGGAGTGGACAGAGAAATTAACAATTACACAAGC

ou on utilise un équivalent codant en conséquence.

11. Procédé selon la revendication 6, caractérisé en ce qu'on utilise en tant que gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA un gène comprenant la séquence de nucléotides suivante :

ATGACGGACAATTGGAGAAGTGAATTATATAAA

TATAAAGTATGTAATAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAGAGAGAGTGGTGCAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTCTGGGTCTTGGGAGCAGCAGGAAGCACTATGGC
 SCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGCTCTGGTATAGTCCAGCAGCAGAACAAT
 TTGCTGACGGCTATTGAGGCGAACAGCATCTGTTCCAACTCAGCTCTGGGGCATCAAGCAGCTCCAG
 GCAAGAACTCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCTCGGGATTTGGGGTTGCTCTCGA
 AACTAATTTCCACCCTCTCTGTCCTTGGAACTGCTAGTTGGAGTAATAAATCTCTGGAACAGATTG
 AATCACACGACGTGGATGGAGTGGACAGAGAAATTAACAATTACACAAGC

ou on utilise un équivalent codant en conséquence.

12. Procédé selon l'une quelconque des revendications 6 à 11, dans lequel le vecteur d'expression est un plasmide pouvant subir une répllication dans des bactéries gram-négatives.

13. Procédé selon la revendication 12, dans lequel le plasmide peut subir une répllication dans une souche de E. coli.

14. Procédé pour préparer un transformant portant un vecteur d'expression, qui comprend un gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA, ce procédé consistant à transformer un micro-organisme avec un vecteur d'expression obtenu selon l'une quelconque des revendications 6 à 13, et à cultiver le micro-organisme transformé.

15. Procédé selon la revendication 14, dans lequel le micro-organisme est une souche de E. coli.

16. Procédé selon la revendication 15, dans lequel le micro-organisme est une souche de E. coli MC 1061.

17. Procédé pour détecter dans le sang humain la présence d'anticorps contre l'agent étiologique viral du SIDA, qui consiste à mélanger une composition contenant un fragment d'une protéine d'enveloppe du virus du SIDA obtenue selon la revendication 1 avec un échantillon de sang humain, et à déterminer si le fragment de la protéine d'enveloppe du SIDA se lie aux anticorps anti-SIDA présents dans l'échantillon sanguin.

18. Procédé selon la revendication 17, qui consiste à utiliser une analyse par "Western Blotting".

19. Procédé selon la revendication 17, qui consiste à utiliser une technique de liaison enzymatique Elisa, dans laquelle un fragment d'une protéine d'enveloppe d'un virus du SIDA obtenue selon la revendication 1 est appliquée sur une phase solide et mise en contact avec l'échantillon et, après lavage, mise en contact avec une IgG non humaine marquée par une enzyme.

20. Procédé selon la revendication 17, dans lequel on utilise la Méthode du Double Antigène .

21. Procédé pour la détermination du virus du SIDA, dans lequel on utilise des anticorps contre un fragment d'une protéine d'enveloppe d'un virus du SIDA obtenue selon la revendication 1.
- 5 22. Procédé selon la revendication 21, dans lequel l'antigène présent dans l'échantillon et un fragment d'une protéine obtenue selon la revendication 1 sous forme marquée entrent en concurrence avec un anticorps contre un fragment d'une protéine obtenue selon la revendication 1.
23. Procédé selon la revendication 21, dans lequel on utilise une méthode sandwich en utilisant deux anticorps contre un fragment d'une protéine obtenue selon la revendication 1.
- 10 24. Procédé selon la revendication 23, dans lequel un anticorps se trouve sur une phase solide et l'autre anticorps est marqué.
25. Procédé selon la revendication 23, dans lequel on utilise deux anticorps monoclonaux différents.
- 15 26. Fragment d'une protéine d'enveloppe d'un virus du SIDA, préparée par un procédé selon l'une quelconque des revendications 1 à 5.
27. Vecteur d'expression comprenant un gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA, préparée par un procédé selon l'une quelconque des revendications 6 à 13.
- 20 28. Transformant portant un vecteur d'expression comprenant un gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA, préparé par un procédé selon l'une quelconque des revendications 14 à 16.
- 25 29. Vecteur d'expression comprenant un gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA selon la revendication 1, en aval d'un promoteur permettant la transcription, la traduction et donc l'expression du fragment de la protéine d'enveloppe dans une cellule hôte.
- 30 30. Vecteur d'expression selon la revendication 29, dans lequel le gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA est un gène comprenant la séquence de nucléotides suivante :

35

40

45

50

55

GTGTGGAAAGGAAGCA

ACCACCCTCTATTTGTGCATCAGATGCTAAAGCATATGATACAGAGGTRCATATTTTGGGCCACA
 CATGCCCTGTGTACCCACAGACCCCAACCCACAAGAAGTAGTATTGGTAAATGTGACAGAAAAATTTAAAC
 ATGTGGAAAAATGACATGGTAGAACAGATGCATGACCATATAATCAGTTTATCGGATCAAGCCCTAAAG
 CCATGTGTAAATTAACCCCACTCTGTGTAGTTTAAAGTGCCTGATTGAAAGATGATACTAATACC
 AATAGTAGTAGCGGGAGAAATGATAATGGAGAAAGGAGAGATAAAAAACTGCTCTTTCAATATCAGCACA
 AGCATAGAGCGTAAGGTGCAGAAAGAATATGCATTTTTTATAAACTTGATATAATACCAATAGATAAT
 GATACTACCACTATACGTTGACAAGTTGTAAACCTCAGTCATTACACAGGCCCTGTCCAAAGGTATCC
 TTTAGCCCAATTCCTATACATTTATGTGCCCCGGCTGGTTTGGGATTCTAAATGTAAATAAAGACC
 TTCAAAGGAACAGGACCATGTACAAAATGTCAGCACAGTACAATGTACACATGGAATTAGGCCAGTAGTA
 TCACTCAACTGCTGTTAAATGGCACTTAGCAAGAAAGAGTAGTAATTAGATCTGTCAATTTACCG
 GACAACTGTAAGCCATAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAAGCCCAACAAAC
 AATACAAAGAAAAAAATCCGTATCCAGAGGGGACCAAGGAGAGCATTGTTTACAATAGCAAAAAATAGGA
 AATATGAGACAAAGCACATTGTAAACATTAGTAGAGCAAAATGGAAATGCCACTTTAAAAACAGATAGCTAGC
 AAATTAAGAGAACAAATTTGGAAATAATAAAACAAATATCTTTAAGCAATCCTCAGGAGCGGACCCAGAA
 ATTGTAAAGCACAGTTTTAATTGTGGAGGGGAATTTTCTACTGTAAATCAACACAATCTGTTAATAGT
 ACTTGGTTTAAATAGTACTTGGAGTACTGAAGGCTCAATAACACTGAGGAGTGCACAAATCAGCTC
 CCATGCAAAATAAAACAAATTTATAAACAATGTGCCAGGAAGTAGGAAAAGCAATGTATGCCCCCTCCCATC
 AGCGACAAATTACATGTTTCAATATATTACAGCGCTGCTATTAAACAGAGATGGTGGTAATACCAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGACAAATTCGAGAAAGTGAAATATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGGCAAGAGAGTGGTGCAGAGA
 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGTTCTTCCGACCAAGCAAGCAACCACTATGGGC
 GCAAGCTCAATCAGGCTGACCGTACAGGCCAGACATTTATGTTCTGTTATAGTGCAGCAGCAGAAAT
 TTGCTGAGCGCTATTGAGGCCCAACAGCATCTGTTGCACTCAGCTCTGGCCATCAAGCAGCTCCAG
 GCAAGAACTCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGATTTGGGTTGCTCTGGA
 AACTAATTTGCACCACTGCTGTGCTTGGAAATGCTAGTTGGAGTAATAAATCTCTGCAACAGATTTGG
 AATCAGACGACGTGGATGGAGTGGACAGAGAAATTAACAAATTACCAAGC

ou l'un de ses équivalents codant pour ledit fragment de ladite protéine d'enveloppe.

31. Vecteur d'expression selon la revendication 29, dans lequel le gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA est un gène comprenant la séquence de nucléotides suivante :

[illegible]

25 ou l'un de ses équivalents codant pour ledit fragment de la protéine d'enveloppe.

32. Vecteur d'expression selon la revendication 29, dans lequel le gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA est un gène comprenant la séquence de nucléotides suivante :

30 ATGAGCAGCAGCATTGTACATTAGTACAGCAGAAATGCAATGCCACTTTAAJACAGATAGCTAGC
AAATTAAGAGAACAAATTTGGAAATAATAAACRAATATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA
ATTGTAACGCACAGCTTTTAAITGTGTGGAGGGGAATTTTCTACTGTAAATTCACACAACCTGTTTAAATAGT
ACTTCGTTTTAATATCTTGTAGTACTGAAGCGGTCAATAACACTGAAGGAAGTGACACAATCACACTC
35 CCATGCAGAAATAAAACRAATTTATAAACATGTUGCAGGAAGTAGGAAAAGCAATGTATGCCCTCCCATC
AGCCGACAAATTAGATGTTTCATCAAAATATTACACCGCTCTCTATTAACAAGAGATGTTGTAATAACAAC
AATCCCTCCGAGATCTTCAAGCCTGGAGGACGAGATATGAGGGACAATTTGGAGAAGTGAATTATATAAA
TATAAAGTAGTAAAAATTGAACCAATTACGAGTAGCACCCACCAAGGCCAAAGAGAAGAGTGGTGACAGAGA
GAAFAAAGAGCAGTGGGAATAGGAACCTTTGTTCTTGGGTTCTTTGGAGCAGCAGGAAGCAATATGGGC
40 GCAGCGTCAATGACCGTGACCGTACAGGCCAGACAATATTGTCTGGTATAGTGCAGCAGCAGAAACAAAT
TTGCTGAGGGCTATTGAGGGCGAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGAGCTCCAG
GCAAGAAATCCTGGCTGTGGAAAGATACCTAAAGCATCAACAGCTCTGCGGATTTGGCGTTGCTCTGGA
AACTAATTTCACCACTGCTGTGCTTGAATGCTAGTTGAGTAATAAAATCTCTGCAACAGATTGGA
AATCACACGAGCTGGATCGAGTGGGACAGAGAAATTAACAAATTACACAAGC

ou l'un des ses équivalents codant pour ledit fragment de la protéine d'enveloppe.

- 50 33. Vecteur d'expression selon la revendication 29, dans lequel le gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA est un gène comprenant la séquence de nucléotides suivante :

ATGTATGCCCCCTCCCATC

5 ACCGGACAAATTAGATGTTTCATCAAATATTACAGCGCTGCTATTAAACAGAGATGGTGGTAATAACAAC
 AATGGGTCCGAGATCTTCAGACCTGGAGGAGGAGATATGAGGACAAATTGGAGAGTGAATTATATAAA
 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGCCAAAGAGAAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATACGAOCTTGTTCCTTGGGTTCCTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACCGTACAGCCAGACAATTATTGTCTGGTATAGTGCAAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGGCGAACACCACTCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 10 GCAAGATCTCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AACTTAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTGGAGTAATAAATCTCTGGAACAGATTGGA
 ATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

ou l'un de ses équivalents codant pour ledit fragment de la protéine d'enveloppe.

34. Vecteur d'expression selon la revendication 29, dans lequel le gène codant pour un fragment d'une protéine d'enveloppe d'un virus du SIDA est un gène comprenant la séquence de nucléotides suivante :

ATGAGGGACAATTGGAGAGTGAATTATATAAA

20 TATAAAGTAGTAAAAATTGAACCATTAGGAGTAGCACCCACCAAGCCAAAGAGAGAGTGGTGCAGAGA
 GAAAAAAGAGCAGTGGGAATACGAGCTTGTTCCTTGGGTTCCTGGGAGCAGCAGGAAGCACTATGGGC
 GCAGCGTCAATGACGCTGACCGTACAGCCAGACAATTATTGTCTGGTATAGTGCAAGCAGCAGAACAAAT
 TTGCTGAGGGCTATTGAGGGCGAACACCACTCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG
 25 GCAAGATCTCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTTGCTCTGGA
 AACTTAATTTGCACCACTGCTGTGCCTTGGAAATGCTAGTGGAGTAATAAATCTCTGGAACAGATTGGA
 ATCACACGACGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGC

35. Vecteur d'expression selon l'une quelconque des revendications 29 à 34, qui est un plasmide pouvant subir une répllication dans des bactéries grain-négatives.

36. Vecteur d'expression selon la revendication 35, qui peut subir une répllication dans une souche de E. coli.

37. Vecteur d'expression pEV1, -2 ou -3/env 44-640.

38. Vecteur d'expression pEV1, -2 ou -3/env 205-640.

39. Transformant portant un vecteur d'expression selon l'une quelconque des revendications 29 à 38.

40. Transformant selon la revendication 39, qui est une souche de E. coli.

41. Transformant selon la revendication 40, qui est une souche de E. coli MC 1061.

42. Anticorps produits contre un fragment d'une protéine obtenue selon les revendications 1 à 5 et 26.

43. Anticorps selon la revendication 42, qui sont des anticorps monoclonaux.

44. Vaccin déclenchant une immunité au SIDA, comprenant comme principe actif un fragment d'une protéine obtenue selon les revendications 1 à 5 et 26.

45. Utilisation d'un fragment d'une protéine selon la revendication 1 pour préparer un vaccin d'immunisation protectrice.

FIGURE 1

1 ATTCGTGCAACAACTGCTGTTTATCCATTTTCAGAAATTOGGTGTGACATAGCAGAATAGGCGTTACTCG 69
 70 ACAGAGGAGAGCAAGAAATGGAGCCAGTAGATCCTAGACTAGAGCCCTGGAGCATCCAAGAGTCAGC 138
 139 CTAAACTGCTTGTACCAATTGCTATGTAAAAAGTGTGCTTTTCATTGCCAAGTTTGTTCATAACAA 207
 208 AAGCCTTAGGCATCTCCTATGGCAGGAAGAGCGAGACAGCGACGAAGACCTCCTCAAGGCAGTCAGA 276
 277 CTCATCAAGTTTCTCTATCAAAGCAGTAAGTAATACATGTAATGCAACCTATACAAATAGCAATAGTAG 345
 346 CATTAGTAGTAGCAATAATAATAGCAATAGTTGTGTGGTCCATAGTAATCATAGAATATAAGGAAAATAT 414
 415 TAAGACAAAGAAAAATAGACAGGTTAATTGATAGACTAATAGAAAGAGCAGAAACAGTCGGCAATGAGA 483
 484 GTGAAGGAGAAATATCAGCACTTGTGGAGATGGGGTGGAGATGGGGCACCATGCTCCTTGGGATGTTG 552
 553 ATGATCTGTAGTGCTACAGAAAAATTTGGGGTCAAGTCTATTATGGGGTACCTGTGTGGAAAGGAGCA 621
 622 ACCACCACTCTATTTTGTGCATCAGATGCTAAAGCATATGATACAGAAGTACATAATGTTTGGGCCACA 690
 691 CATGCTGTGTACCCACAGACCCCAACCCCAAGAGAGTAGTATTGTTAAATGTGACAGAAAAATTTTAAC 759
 760 ATGTGAAAAATGACATGTGAGAACAGATGATGAGATATAATCAGTTTATGGGATCAAAGCCTAAAG 828
 829 CCATGTGTAAAATTAACCCCACTCTGTGTTAGTTTAAAGTGCAGTATTTGAGAAATGATACTAATACC 897
 898 AATAGTAGTAGCGGAGAAATGATAATGGAGAAAGAGAGATAAAAAACTGCTCTTTCAATATCAGCACA 966
 967 AGCATAAGAGGTAAAGTGCAGAAAGAAATATGCAATTTTTTATAAACTTGATATAATACCAATAGATAAT 1035
 1036 GATACTACCAGCTATACGTTGACAAAGTTGTAACACCTCAGTCATTACACAAGCCCTGTCCAAAGGTATCC 1104
 1105 TTTGAGCCAAATTCOCATACATTATTGTGCCCCCGCTGGTTTGGGATTCTAAAATGTAATAAAGAGC 1173
 1174 TTCAATGGAAACAGACCATGTACAAATGTGACACAGTACAAATGTACACATGGAAATAGGCCAGTAGTA 1242
 1243 TCAACTCAACTGCTGTTAAATGGCAGTCTAGCAGAAAGAGAGTAGTAATTAGATCTGTCAATTTTCAAG 1311
 1312 GACAATGCTAAAACCATTAATAGTACAGCTGAACACATCTGTAGAAATTAATTGTACAAAGACCAACAA 1380
 1381 AATACAAGAAAAAAATTCGGTATCCAGAGGGAGACCAAGGAGAGCATTTGTTACAAAGAAAAATAGGA 1449
 1450 AATATGAGACAAGCACATTGTAACATTAGTAGAGCAAAATGGAAATGCCACTTTAAACAGATAGCTAGC 1518
 1519 AAATTAAGAGAACAATTTGGAAATAATAAAACAATAATCTTTAAGCAATCCTCAGGAGGGGACCCAGAA 1587
 1588 ATTGTAAACACAGTTTAAATTTGTGGAGGGGAAATTTTCTACTGTAATTCAACACAAGTGTTAATAGT 1656
 1657 ACTTGGTTTAATAGTACTTGGAGTACTGAAGGGTCAAAATACACTGAAGGAAAGTGACACAATCACACTC 1725
 1726 CCATGCAGAATAAAACAATTTATAACATGTGGCAGGAAGTAAGAAAAGCAATGTATGCCCTCCCATC 1794
 1795 AGCGGACAAAATTAGATGTTCAATCAAAATATTACAAGGCTGCTATTAAACAAGAGATGGTGGTAATAACAA 1863
 1864 AATGGGTCGAGATCTTCAGACCTGGAGGAGAGATATGAGGACAATTGGAGAAGTGAATTATATAAA 1932
 1933 TATAAGTAGTAAAAATTGAACCATTAAGAGTAGCAACCCACCAAGGCAAGAGAGAGTGGTGACAGAGA 2001
 2002 GAAAAAGAGCAGTGGGAATAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTATGGGC 2070
 2071 GCAGCGTCAATGAGCGTGACGGTACAGGCCAGACAATTAATTGTCTGGTATAGTGACAGCAGCAGAACAA 2139
 2140 TTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAG 2208
 2209 GCAAGAACTCTGGCTGTGGAAGATACCTAAAGGATCAACAGCTCCTGGGGATTGGGGTGTCTCTGGA 2277
 2278 AAATAATTGACCACTGCTGTGCTTGGAAATCTAGTGGAGTAATAAATCTCTGGAACAGATTGAG 2346
 2347 AATCACACAGCGTGGATGGAGTGGGACAGAGAAATTAACAATTACACAAGCTTAATACACTCTTAATT 2415
 2416 GAAGAATCGCAAAACCAAGCAAGAAAGAAATGAACAGAAATTAATGAAATTAATGAAATGGCAAGTTG 2484
 2485 TGAATTTGTTTAAACATAACAAATGGCTGTGGTATATAAAATTTTCAATAATGATAGTAAGGAGCTT 2553
 2554 GTAGGTTTAAGAAATAGTTTGTGCTGACTTTCTGTAGTGAATAGAGTTAGCCAGGGATATTCACCATTA 2622
 2623 TCGTTTCAGACCCACCTCCCAATCCCGAGGGGACCCGACAGGCCGGAAGAAATAGAAAGAAAGGTGGA 2691
 2692 GAGAGAGACAGAGACAGATCCATTGAGATTAGTGAACGATTCCTTAGCACTTATCTGGGACGATCTGGGG 2760
 2761 AGCCTGTGCTCTTCAGCTACCAACGCTTGAGAGACTTACTCTTGATTGTAACGAGGATTGTGAACTT 2829
 2830 CTGGACGCAAGGGGTGGAGCCCTCAAAATATTGTTGAAATCTCTACAAATATTGGAGTCAAGAGCTA 2898
 2899 AAGAATAGTGCTGTAGCTTGGCTCAATGCCACAGCTATAACAGTAGCTGAGGGGACAGATAAGGTTATA 2967
 2968 GAAGTAGTACAAGAGCTTATAGAGCTATTGCCACATACCTAGAAGAAATAGACAGGGCTTGGAAAGG 3036
 3037 ATTTTGCTATAAGATGGTGGCAAGTGGTCAAAAGTAATGTGGTTGGATGGCCTGCTGTAGGGAGAAAG 3105
 3106 AATGAGACGAGCTGAGCCAGCAGCAGATGGGGTGGAGCAAGCATCTCGAGA 3156

FIGURE 2 (3 pages)

| | | | |
|-------|---|----------------|-----------------|
| | 1 | | 50 |
| HXB-3 | MRVKEK-----YQHLWRWGWRWGTMLLGMLMICSATEKLWVTVYYGVPVWKEATT | | |
| BH-10 | | | |
| BH-8 | | | F |
| LAV | | K | I |
| ARV-2 | K --GTRRN | ----- | -- |
| | 51 | | 100 |
| HXB-3 | TLFCASDAKAYDTEVHNVWATHACVPTDPNPQEVVLNVVTENFNMWKNDM | | |
| BH-10 | | | |
| BH-8 | | | |
| LAV | | | |
| ARV-2 | R | G | N |
| | 101 | | 150 |
| HXB-3 | VEQMHEDIISLWDQSLKPCVKLTPLCVSLKCTDLKNDTNTNSS-----SGRMIME | | |
| BH-10 | | | |
| BH-8 | | | |
| LAV | | G A | NTNSS E M |
| ARV-2 | Q | T N | G A NWKEEI----- |
| | 151 | | 200 |
| HXB-3 | KGEIKNCSFNISTSIRGKVQKEYAFFYKLDIIPIDND--TTSYTLTS---CNTSV | | |
| BH-10 | | | |
| BH-8 | K | | |
| LAV | | | |
| ARV-2 | T | D I N L R N VV | AST N NYRLIH R |
| | 201 | | 250 |
| HXB-3 | ITQACPKVSFEPPIPIHYCAPAGFAILKCNNKTFNGTGPCTNVSTVQCTHG | | |
| BH-10 | | | |
| BH-8 | | | |
| LAV | | A | |
| ARV-2 | T | K | |
| | 251 | | 300 |
| HXB-3 | IRPVVSTQLLNGLSLAEVIRSVNFTDNAKTIIVQLNTSVEINCTRPN | | |
| BH-10 | | A | Q |
| BH-8 | | | D |
| LAV | | A | Q |
| ARV-2 | I | D N | E A |

301

350

HXB-3 NNTRKKIRIQRGPGRAFVTIGKIGNMRQ-AHCNISRAKWNATLKQIASKLR
 BH-10 S N D
 BH-8 D
 LAV S
 ARV-2 S Y -- H T R I G D I R K Q N E V K

351

400

HXB-3 EQFGNNKTIIFKQSSGGDPEIVTHSFNCGGEFFYCNSTQLFNSTWPNSTW
 BH-10
 BH-8
 LAV
 ARV-2 V N M R T N -RLNH

401

450

HXB-3 STEGSNNTEGSDTITLPCRICKFINMWQEVGKAMYAPPISGQIRCSSNIT
 BH-10 K I
 BH-8 K I
 LAV
 ARV-2 - --- K N I I G S

451

500

HXB-3 GLLLTRDGG-NNNNGSEIFRPGGGDMRDNRSELYKYKVVKIEPLGVAPTK
 BH-10 - S E
 BH-8 - S E
 LAV -
 ARV-2 T V T D T V I I

501

550

HXB-3 AKRRVVQREKRAVGI-GALFLGFLGAAGSTMGAASMTLTVQARQLLSGIVQ
 BH-10 -
 BH-8 -
 LAV - R
 ARV-2 V M V L

551

600

HXB-3 QQNNLLRAIEAQQHLLQLTVWGIKQLQARILAVERYLKDQQLLGWGC SG
 BH-10 G
 BH-8
 LAV
 ARV-2 V R

601

650

HXB-3 KLICTTAVPWNASWSNKSLEQIWNHTTWMEWDREINNYTSLIHSLIEESQ
 BH-10 NM
 BH-8 NM
 LAV NM
 ARV-2 D DNM Q E D NT YT

651

700

HXB-3 NQOEKNEQELLELDKWASLWNWPNITNWLWYIKLFIMIVGGLVGLRIVFA
 BH-10
 BH-8
 LAV I
 ARV-2 S I

701

750

HXB-3 VLSVNVNRVQGYSPSPQTHLPPIRGPDREPEGIEEGGERDRDRSIRLVN
 BH-10
 BH-8 I N
 LAV I T
 ARV-2 I R V D V D

751

800

HXB-3 GSLALIWDRLRSLCLPSYHRLRDLILLIVTRIVELLGREGWEALKYWNLL
 BH-10
 BH-8
 LAV
 ARV-2 F E R AA T I H S

801

850

HXB-3 QYWSQELKNSAVSLLNATAIAVAEGTDRVIEVVQAYRAIRHIPRRIRQG
 BH-10 G
 BH-8 N L A
 LAV G C
 ARV-2 I W T A R L H

851 856

HXB-3 LERILL
 BH-10
 BH-8
 LAV
 ARV-2 L

" - " designates a deletion of one amino acid. An empty space denotes identity with HXB-3 sequence.

Figure 3




Figure 4



Figure 5





FIGURE 6A

METArg

ValLysGluLysTyrGlnHisLeuTrpArgTrpGlyTrpArgTrpGlyThrMETLeuLeuGlyMETLeu
 METIleCysSerAlaThrGluLysLeuTrpValThrValTyrTyrGlyValProValTrpLysGluAla
 ThrThrThrLeuPheCysAlaSerAspAlaLysAlaTyrAspThrGluValHisAsnValTrpAlaThr
 HisAlaCysValProThrAspProAsnProGlnGluValValLeuValAsnValThrGluAsnPheAsn
 METTrpLysAsnAspMETValGluGlnMETHisGluAspIleIleSerLeuTrpAspGlnSerLeuLys
 ProCysValLysLeuThrProLeuCysValSerLeuLysCysThrAspLeuLysAsnAspThrAsnThr
 AsnSerSerSerGlyArgMETIleMETGluLysGlyGluIleLysAsnCysSerPheAsnIleSerThr
 SerIleArgGlyLysValGlnLysGluTyrAlaPhePheTyrLysLeuAspIleIleProIleAspAsn
 AspThrThrSerTyrThrLeuThrSerCysAsnThrSerValIleThrGlnAlaCysProLysValSer
 PheGluProIleProIleHisTyrCysAlaProAlaGlyPheAlaIleLeuLysCysAsnAsnLysThr
 PheAsnGlyThrGlyProCysThrAsnValSerThrValGlnCysThrHisGlyIleArgProValVal
 SerThrGlnLeuLeuLeuAsnGlySerLeuAlaGluGluGluValValIleArgSerValAsnPheThr
 AspAsnAlaLysThrIleIleValGlnLeuAsnThrSerValGluIleAsnCysThrArgProAsnAsn
 AsnThrArgLysLysIleArgIleGlnArgGlyProGlyArgAlaPheValThrIleGlyLysIleGly
 AsnMETArgGlnAlaHisCysAsnIleSerArgAlaLysTrpAsnAlaThrLeuLysGlnIleAlaSer
 LysLeuArgGluGlnPheGlyAsnAsnLysThrIleIlePheLysGlnSerSerGlyGlyAspProGlu
 IleValThrHisSerPheAsnCysGlyGlyGluPhePheTyrCysAsnSerThrGlnLeuPheAsnSer
 ThrTrpPheAsnSerThrTrpSerThrGluGlySerAsnAsnThrGluGlySerAspThrIleThrLeu
 ProCysArgIleLysGlnPheIleAsnMETTrpGlnGluValGlyLysAlaMETIleTyrAlaProProIle
 SerGlyGlnIleArgCysSerSerAsnIleThrGlyLeuLeuLeuThrArgAspGlyGlyAsnAsnAsn
 AsnGlySerGluIlePheArgProGlyGlyGlyAspMETArgAspAsnTrpArgSerGluLeuTyrLys
 TyrLysValValLysIleGluProLeuGlyValAlaProThrLysAlaLysArgArgValValGlnArg
 GluLysArgAlaValGlyIleGlyAlaLeuPheLeuGlyPheLeuGlyAlaAlaGlySerThrMETGly
 AlaAlaSerMETThrLeuThrValGlnAlaArgGlnLeuLeuSerGlyIleValGlnGlnGlnAsnAsn
 LeuLeuArgAlaIleGluAlaGlnGlnHisLeuLeuGlnLeuThrValTrpGlyIleLysGlnLeuGln
 AlaArgIleLeuAlaValGluArgTyrLeuLysAspGlnGlnLeuLeuGlyIleTrpGlyCysSerGly
 LysLeuIleCysThrThrAlaValProTrpAsnAlaSerTrpSerAsnLysSerLeuGluGlnIleTrp
 AsnHisThrThrTrpMETGluTrpAspArgGluIleAsnAsnTyrThrSerLeuIleHisSerLeuIle
 GluGluSerGlnAsnGlnGlnGluLysAsnGluGlnGluLeuLeuGluLeuAspLysTrpAlaSerLeu
 TrpAsnTrpPheAsnIleThrAsnTrpLeuTrpTyrIleLysLeuPheIleMETIleValGlyGlyLeu
 ValGlyLeuArgIleValPheAlaValLeuSerValValAsnArgValArgGlnGlyTyrSerProLeu
 SerPheGlnThrHisLeuProIleProArgGlyProAspArgProGluGlyIleGluGluGluGlyGly
 GluArgAspArgAspArgSerIleArgLeuValAsnGlySerLeuAlaLeuIleTrpAspAspLeuArg
 SerLeuCysLeuPheSerTyrHisArgLeuArgAspLeuLeuLeuIleValThrArgIleValGluLeu
 LeuGlyArgArgGlyTrpGluAlaLeuLysTyrTrpTrpAsnLeuLeuGlnTyrTrpSerGlnGluLeu
 LysAsnSerAlaValSerLeuLeuAsnAlaThrAlaIleAlaValAlaGluGlyThrAspArgValIle
 GluValValGlnGluAlaTyrArgAlaIleArgHisIleProArgArgIleArgGlnGlyLeuGluArg
 IleLeuLeu

FIGURE 6BAMINO ACID DISTRIBUTION
OF AIDS ENV PROTEIN

| <u>Name</u> | <u>Number of Residues</u> |
|----------------------------|---------------------------|
| A Alanine | 47 |
| B Aspartic Acid-Asparagine | 0 |
| C Cysteine | 21 |
| D Aspartic Acid | 27 |
| E Glutamic Acid | 49 |
| F Phenylalanine | 26 |
| G Glycine | 58 |
| H Histidine | 14 |
| I Isoleucine | 63 |
| K Lysine | 44 |
| L Leucine | 83 |
| M Methionine | 17 |
| N Asparagine | 60 |
| P Proline | 29 |
| Q Glutamine | 42 |
| R Arginine | 52 |
| S Serine | 57 |
| T Threonine | 60 |
| V Valine | 56 |
| W Tryptophan | 31 |
| Y Tyrosine | 20 |
| Z Glutamine-Glutamic Acid | 0 |

Figure 7




Figure 8

